



## FEP Medical Policy Manual

### FEP 8.01.64 Home Non-Invasive Positive Airway Pressure Devices for the Treatment of Respiratory Insufficiency and Failure

**Annual Effective Policy Date: July 1, 2024**

**Original Policy Date: April 2022**

#### **Related Policies:**

2.01.18 - Diagnosis of Obstructive Sleep Apnea Syndrome

8.01.67 - Medical Management of Obstructive Sleep Apnea Syndrome

## Home Non-Invasive Positive Airway Pressure Devices for the Treatment of Respiratory Insufficiency and Failure

### Description

#### Description

Respiratory failure is characterized by low arterial blood oxygen (hypoxemia,  $\text{PaO}_2$ ) and/or high arterial carbon dioxide (hypercapnia,  $\text{PaCO}_2 >45$  mmHg). Chronic respiratory insufficiency or failure can occur with chronic obstructive pulmonary disease (COPD), thoracic restrictive disorders, and hypoventilation syndromes, and may result in poor quality of life, sleepiness, hospital admission, intubation, and death. Non-invasive positive airway pressure ventilation (NPPV) including continuous positive airway pressure (CPAP), bilevel positive airway pressure (BPAP), and home mechanical ventilators (HMV) that are pressure, rate and volume targeted are proposed for the treatment of COPD and other forms of chronic hypoventilation.

#### Chronic Obstructive Pulmonary Disease

Chronic obstructive pulmonary disease (COPD) is a common condition, affecting more than 5% of the population, and is associated with high morbidity and mortality. COPD is the fourth leading cause of death in the United States. It is a clinical syndrome with multiple etiologies that is characterized by chronic respiratory symptoms, structural pulmonary abnormalities, and/or lung function impairment. Chronic obstructive pulmonary disease is most frequently associated with cigarette smoking or other air pollutants, and a majority of patients with COPD in the United States have a history of cigarette smoking. Chronic obstructive pulmonary disease is progressive, with expiratory airflow limitation, air trapping/hyperinflation, and destruction of alveoli (emphysema). The Global Initiative for Chronic Obstructive Lung Disease (GOLD), defines COPD as "a heterogeneous lung condition characterized by chronic respiratory symptoms (dyspnea, cough, sputum production and/or exacerbations) due to abnormalities of the airways (bronchitis, bronchiolitis) and/or alveoli (emphysema) that cause persistent, often progressive, airflow obstruction".<sup>1,2</sup>

Respiratory failure in patients with COPD is characterized by the inability to sustain normal gas exchange, leading to low arterial blood oxygen (hypoxemia, PaO<sub>2</sub>) and/or high arterial carbon dioxide (hypercapnia, PaCO<sub>2</sub>). Hypercapnia develops in about one-third of patients with COPD and is associated with poor quality of life, sleepiness, frequent hospital admissions due to exacerbations, and an increase in mortality compared to patients with COPD who are normocapnic. The hypercapnia is due in large part to poor lung biomechanics including low inspiratory muscle reserve, high CO<sub>2</sub> production, and a reduced ventilatory capability.<sup>3</sup> The imbalance between the respiratory load and respiratory capability may in turn affect the ventilatory control center in the brain stem. Physiological changes in responsiveness to hypoxemia and hypercapnia during sleep can be particularly pronounced in patients with COPD, with overnight increases in PaCO<sub>2</sub> affecting daytime PaCO<sub>2</sub>, possibly through bicarbonate retention or changes in cerebrospinal fluid.<sup>4</sup> Patients with COPD may also have comorbid obstructive sleep apnea and/or obesity hypoventilation syndrome due to decreased ventilatory motor output and upper airway muscle activity during sleep.

## Thoracic Restrictive Disorders Due to Neuromuscular Disease

Thoracic restrictive disorders result from a variety of underlying diseases all characterized by restrictive patterns on pulmonary function testing.<sup>5</sup> Neuromuscular disorders such as muscular dystrophy, amyotrophic lateral sclerosis (ALS), polio, and phrenic neuropathies can result in weakness of the respiratory muscles affecting inspiration and expiration, ultimately resulting in hypoventilation. Impaired cough and swallowing associated with neuromuscular disease increases the risk of respiratory complications in these patients.<sup>6</sup> Nocturnal hypoventilation due to muscular atonia during sleep leads to nocturnal hypercapnia. Frequent nocturnal episodes can result in renal compensation and ultimately result in daytime hypercapnia. Non-invasive positive airway pressure ventilation (NPPV) is often necessary for patients with thoracic restrictive disorders due to neuromuscular disease.

## Hypoventilation Syndromes

Hypoventilation syndromes are nonspecific disorders characterized by hypercapnia (PaCO<sub>2</sub> >45 mmHg) that is not otherwise categorized.<sup>7</sup> Obesity hypoventilation syndrome (OHS), central respiratory depression due to substance or medication use, and decompensated hypercapnic respiratory failure that is not COPD are all included in this category. In patients with OHS, weight loss is useful in normalizing PaCO<sub>2</sub>; however, NPPV should be initiated early while weight loss is attempted.<sup>8</sup>

## Treatment With Non-invasive Positive Airway Pressure

A major goal of management of patients with chronic hypoventilation is to reduce hospitalizations and mortality. Long-term oxygen therapy is recommended for patients with poor clinical status and NPPV devices for patients with severe chronic hypercapnia and a history of hospitalization for acute respiratory failure. Non-invasive positive airway pressure ventilation devices include nocturnal continuous positive airway pressure (CPAP) for individuals with hypercapnia due to obstructive sleep apnea or hypoventilation and bilevel positive airway pressure (BPAP) devices or non-invasive home mechanical ventilators that are pressure, rate, and volume targeted. The objective of this evidence review is to describe which features of NPPV are required to improve the net health outcome in patients with COPD, thoracic restrictive disorders due to neuromuscular disease, or those with hypoventilation syndromes.

Benefits of nocturnal NPPV persist into the daytime with improved breathing patterns (lower frequencies and larger tidal volumes) and improved gas exchange. Explanations for the improvement in daytime respiration with nocturnal NPPV include increased respiratory drive, improved diaphragm function by unloading the respiratory muscles during sleep, increased CO<sub>2</sub> sensitivity, and reduction in air trapping and hyperinflation. It is not known which factors (eg, muscle unloading, gas exchange normalization, decrease in hyperinflation) underlie the benefits of NPPV on health outcomes. It is also unclear if the reduction in PaCO<sub>2</sub> has an effect on health outcomes or if it is only a marker of effective ventilation.<sup>4</sup>

## Respiratory Assist Devices

The Centers for Medicare and Medicaid Services (CMS) defines respiratory assist devices (RADs) as bilevel devices with or without back-up respiratory rate capability. While CPAP devices provide continuous air at a pressure that prevents the collapse of the airway during inspiration, BPAP devices work by increasing pressure during inspiration and lowering it during expiration (pressure cycled). In some devices a backup respiratory rate is triggered when the patient's nocturnal respiratory rate decreases below a set threshold. The backup rate is typically set 2 breaths below the patient's spontaneous respiratory rate during wakefulness.

Terminology on device features is described in Table 1.

### Table 1. Device Features

Term	Definition	Description
Bilevel-S	Bilevel without a backup rate	Positive airway pressure that is higher during inspiration than expiration that is triggered by patient inspiration.
Bilevel-ST	Bilevel with a backup rate	Positive airway pressure that is higher during inspiration than expiration with a backup respiratory cycle length if the patient's breathing is slower than the preset rate.
VAPS	Volume-assured pressure support modes	Bilevel ST modes that use an algorithm to adjust inspiratory pressure support to meet a set tidal volume.
iVAPS	Intelligent volume-assured pressure support modes	Bilevel ST modes that use an algorithm to adjust inspiratory pressure support within a predetermined range to meet a set target ventilation.

## Home Mechanical Ventilators

In some patients, nocturnal respiratory assist devices are insufficient to address the respiratory failure. Non-invasive home mechanical ventilators (HMV) are proposed for the treatment of chronic respiratory failure that is refractory to a respiratory assist device. Mechanical ventilators are devices that deliver more controlled breathing with bilevel ventilation at a higher pressure. The ventilators may also have additional features compared to BPAP machines such as alarms and battery backup power. Home mechanical ventilators can be used for patients with tracheostomy in the home, but may also be used with a non-invasive interface such as a mask or mouthpiece in patients who do not depend on 24 hour ventilation for survival. Current technology has decreased the size of home ventilators to around 10 pounds. In addition, some models may be wireless with battery backup, allowing greater mobility during the day.

## Titration

Early studies with low intensity NPPV did not demonstrate health benefits in patients with hypercapnia. More recent studies have reinforced the importance of high-intensity NPPV (>18 cm H<sub>2</sub>O) that is titrated to decrease hypercapnia. A high respiratory backup rate that is increased to the level of spontaneous breathing has also been shown to be important to achieve positive health outcomes. Manually set, laboratory or hospital titration of NPPV with pressure control and backup rate have been recommended for stable hypercapnic COPD.<sup>9</sup> The goal of titration of inspiratory positive airway pressure is to achieve normocapnia, a reduction in transcutaneous CO<sub>2</sub>, or maximum tolerable inspiratory pressure. A fast rise in inspiratory pressure (rise time) allows enough time for expiration within the normal rate of breathing. In patients with air trapping and hyperinflation, use of positive end-expiratory pressure can also be beneficial.

A suggested protocol for in-laboratory titration of NPPV in patients with COPD in the U.S. is described by Orr et al (2020).<sup>4</sup> Titration of NPPV is usually performed in a monitored environment after the patient has stabilized, as studies have not found an improvement in health outcomes when NPPV is started soon after an acute exacerbation. Polysomnography or respiratory monitoring may be used during titration to evaluate the presence of obstructive sleep apnea or hypoventilation. The inspiratory pressure is typically started at 6 to 8 cm H<sub>2</sub>O of pressure support above the expiratory pressure and titrated to reduce hypercapnia. A Bilevel-ST (with backup rate) or a VAPS (volume assured) may be used if a Bilevel-S (without backup rate) fails to adequately reduce hypercapnia. Although titration in European studies has been performed with a hospital stay, this is not feasible in the U.S., and titration might be performed over several weeks in the patient's home by an external durable medical equipment (DME) provider.

## Pulmonary Rehabilitation

Pulmonary rehabilitation is a personalized intervention that includes physical activity (eg, activities of daily living, endurance exercises and muscle strengthening), health education, and psychological support. It may be performed in the hospital, outpatient clinic, or home, and has been shown to reduce mortality, exacerbation rate, intensive care admissions, and emergency department visits. Pulmonary rehabilitation is common in Europe but is less frequently provided in the U.S.

## OBJECTIVE

The objective of this evidence review is to determine which features of non-invasive positive airway pressure improve the net health outcome in individuals with COPD, thoracic restrictive disorders due to neuromuscular disease, or hypoventilation syndromes including obesity hypoventilation syndrome.

## POLICY STATEMENT

For individuals with chronic obstructive pulmonary disease (COPD) without significant hypercapnia ( $\text{PaCO}_2 < 52$  mmHg) and with obstructive sleep apnea, see policy 2.01.18 (Diagnosis of Obstructive Sleep Apnea Syndrome) and 8.01.67 (Medical Management of Obstructive Sleep Apnea Syndrome).

Nocturnal bilevel positive airway pressure (BPAP) with backup rate may be considered **medically necessary** for individuals with COPD and chronic respiratory failure (see Policy Guidelines) who meet either of the following:

- Chronic stable daytime (awake) hypercapnia ( $\text{PaCO}_2 > 52$  mmHg) **OR**
- Daytime (awake) hypercapnia ( $\text{PaCO}_2 > 52$  mmHg) at least 2 weeks after discharge from the hospital for an acute exacerbation with decompensated respiratory acidosis.

Non-invasive home mechanical ventilation may be considered **medically necessary** for individuals with COPD who meet the following criteria:

- Qualify for a BPAP device **AND** meet at least one of the following:
  - Higher pressure is needed to reduce hypercapnia than can be achieved with a bilevel device during titration (typically  $> 25$  cm  $\text{H}_2\text{O}$ ); **OR**
  - Severe hypoxemia requiring fraction of inspired oxygen ( $\text{FIO}_2$ )  $> 40\%$  or  $> 5$  L/min; **OR**
  - Daytime use (battery operated unit) is required to reduce hypercapnia.

Individuals with COPD who are started on BPAP at discharge from hospitalization may continue for up to 3 months to provide time to stabilize and complete reevaluation.

Non-invasive positive airway pressure for COPD is considered **investigational** under all other conditions.

Bilevel positive airway pressure may be considered **medically necessary** for individuals with thoracic restrictive disorders due to neuromuscular disease who meet any of the following:

- Pulmonary function tests:
  - Spirometry (upright or supine) with vital capacity  $< 50\%$  predicted or  $< 80\%$  predicted with associated symptoms (orthopnea, dyspnea, morning headaches, excessive daytime sleepiness, or unrefreshing sleep); **OR**
  - Maximal inspiratory pressure  $< 60$  cm  $\text{H}_2\text{O}$  or maximum expiratory pressure (MEP)  $< 40$  cm  $\text{H}_2\text{O}$ ; **OR**
  - Peak cough flow (PCF)  $< 270$  L/min for age  $\geq 12$  years or PCF  $< 5$ th percentile for age  $< 12$  years; **OR**
  - Sniff nasal inspiratory pressure (SNIP)  $< 70$  cm  $\text{H}_2\text{O}$  in males, SNIP  $< 60$  cm $\text{H}_2\text{O}$  in females for age  $\geq 12$  years.
- Hypercapnia
  - Chronic stable daytime (awake) hypercapnia with  $\text{PaCO}_2 \geq 45$  mmHg (capillary blood gas can be used in children); **OR**
  - Venous blood gas  $\text{PCO}_2$ , end-tidal  $\text{PCO}_2$ , or transcutaneous  $\text{PCO}_2$ ,  $\geq 50$  mmHg; **OR**
- Hypoxia
  - Overnight oximetry in-laboratory or home sleep test with saturation  $< 88\%$  for 5 minutes;
  - Overnight oximetry:  $\text{SpO}_2 \leq 90\%$  for  $\geq 2\%$  of sleep time.

Non-invasive home mechanical ventilation may be considered **medically necessary** for individuals with thoracic restrictive disorders due to neuromuscular disease who meet the following:

- Qualify for a BPAP device; **AND**

- o BPAP fails; OR
- o Have extreme loss in function with vital capacity <30%; OR
- o Non-invasive ventilation is needed for >10 hours per day; OR
- o Severe breathlessness (e.g., with speaking at rest); OR
- o Worsening daytime hypercapnia with need for mouthpiece ventilation; OR
- o Daytime use (battery operated unit) is required to reduce hypercapnia or dyspnea.

Bilevel positive airway pressure may be considered **medically necessary** for individuals with hypoventilation syndromes who meet the following criteria:

- Awake or sleep hypoventilation with hypercapnia (one of the following is met):
  - o Awake hypoventilation with chronic stable daytime (awake) hypercapnia ( $\text{PaCO}_2 \geq 45$  mmHg); OR
  - o Venous blood gas  $\text{PCO}_2$ , end-tidal  $\text{PCO}_2$ , or transcutaneous  $\text{PCO}_2 \geq 50$  mmHg; OR
  - o Sleep hypoventilation with hypercapnia:
    - $\geq 10$  mmHg increase from baseline awake  $\text{PCO}_2$  and to a value  $> 50$  mmHg for  $\geq 10$  min; OR
    - $\text{PCO}_2 \geq 55$  mmHg for  $\geq 10$  min; AND
- Low clinical suspicion for COPD or neuromuscular disease; AND
- One of the following conditions are met:
  - o Obesity with body mass index (BMI)  $\geq 30$   $\text{kg/m}^2$ ; OR
  - o Decreased respiratory drive due to opioid or substance use; OR
  - o Advanced lung disease other than COPD (e.g., end-stage or advanced interstitial lung disease); AND
- Individual was discharged from inpatient stay with persistent awake hypoventilation (hypercapnia) on BPAP.
  - o A reassessment with a provider within 3 months (30 to 90 days) is required and an attended polysomnogram (PSG) should be performed to assess appropriateness of positive airway pressure modality (home sleep apnea test is acceptable if attended PSG is not obtainable); OR
- Individual is ambulatory and sleep study indicates that BPAP is necessary for sleep-disordered breathing, or individual with severe obstructive sleep apnea is continuous positive airway pressure intolerant or continuous positive airway pressure was proven ineffective.

Non-invasive home mechanical ventilation may be considered **medically necessary** for individuals with hypoventilation syndromes who meet the following:

- Qualify for a BPAP device and at least one of the following:
  - o Higher pressure is needed to reduce hypercapnia than can be achieved with a bilevel continuous positive airway pressure device during titration (typically  $>25$  cm  $\text{H}_2\text{O}$ ); OR
  - o Severe hypoxemia requiring  $\text{FIO}_2 >40\%$  or  $>5$  L/min; OR
  - o Daytime use (battery operated unit) is required to reduce hypercapnia; OR
- Tried and failed BPAP device with persistent hypercapnia despite 3 months of adequate adherence to prescribed positive airway pressure therapy with:
  - o Awake  $\text{PaCO}_2 \geq 45$  mmHg; OR
  - o Awake venous blood gas  $\text{PCO}_2$ , end-tidal  $\text{PCO}_2$ , or transcutaneous  $\text{PCO}_2 \geq 50$  mmHg.

## POLICY GUIDELINES

Respiratory failure in individuals with chronic disease is characterized by the inability to sustain normal gas exchange, leading to low arterial blood oxygen (hypoxemia, PaO<sub>2</sub>) and/or high arterial carbon dioxide (hypercapnia, PaCO<sub>2</sub>). Assessment of hypoxemia would lead to supplemental oxygen administration. Stable clinical state is defined as free of exacerbations for at least 4 weeks with pH over 7.35.

Compliance with treatment of at least 4 hours per 24 hours should be documented after the first 3 months of use. There are limited data on which to base compliance assessment. Assessment could be further based on an *average* of at least 4 hours per 24 hours over a consecutive 30-day period or use of 4 hours per 24 hours for at least 65% of the days in a consecutive 30-day period.

The Centers for Medicare and Medicaid Services (CMS) classifies a respiratory assist device as a bilevel positive airway pressure (BPAP) device with or without backup respiratory rate capability. Treatment modalities that are reported with the E0471 code include BiPAP ST, ASV, BiPAP AutoSV, iVAPS, AVAPS. BPAP units with batteries have a battery life that is shorter than home mechanical ventilators and are infrequently used in the U.S.

CMS defines non-invasive mechanical ventilators as life supporting/sustaining devices used in various settings, including home, hospital, and institutional settings. The non-invasive mechanical ventilators should have at least 6 pressure modes and 3 volume modes, and allow for both invasive or non-invasive use. For examples, see the Regulatory Status section.

Although most patients with comorbid COPD and obstructive sleep apnea can be effectively treated with continuous or auto-adjusting positive airway pressure, approximately 10% of patients will need BPAP to tolerate the required pressure. These devices are reviewed in reference medical policy 2.01.18 (Diagnosis of Obstructive Sleep Apnea Syndrome) and 8.01.64 (Medical Management of Obstructive Sleep Apnea Syndrome).

Respiratory therapy in the home may be provided for patients who are treated with E0466, E0470, or E0471 devices.

## BENEFIT APPLICATION

Experimental or investigational procedures, treatments, drugs, or devices are not covered (See General Exclusion Section of brochure).

## FDA REGULATORY STATUS

Numerous CPAP and BPAP devices are available in the U.S. Examples of HMV devices that have both invasive and non-invasive interfaces and are available in the U.S. are described in Table 2.

**Table 2. Select Home Mechanical Ventilators with Non-invasive Interface**

Device	Manufacturer	FDA clearance	Date	FDA product code
Trilogy™ Evo Ventilator	Respironics	K181166	2019	NOU, CBK
Vivo 60	Breas	K160481	2016	NOU, CBK, DQA, CCK
Astral 100/150	ResMed	K152068	2016	NOU, CBK
Newport™	Medtronic	K121891	2012	NOU, CBK
iVent	GE Healthcare	K092135	2009	NOU, CBK
LTV	Cardinal Health	K083688	2009	CBK



Puritan Bennett 540

Covidien

K082966

2008

CBK

## RATIONALE

### Summary of Evidence

For individuals who have chronic obstructive pulmonary disease (COPD) and obstructive sleep apnea (OSA) who receive continuous positive airway pressure (CPAP), the evidence includes observational studies. Relevant outcomes are mortality, symptoms, morbid events, functional outcomes, quality of life, and hospitalization. Studies of patients with both COPD and OSA who do or do not use CPAP show a mortality benefit in patients with overlap syndrome who are treated with positive airway pressure. The greatest benefits occur in patients with COPD and hypercapnia and in older adults, and individuals with more comorbid conditions and higher complexity ratings. It should be noted that the threshold for what was considered hypercapnia was lower than in other studies on bilevel positive airway pressure (BPAP) that used a threshold of arterial blood carbon dioxide ( $\text{PaCO}_2$ )  $>52$  mmHg. Although the literature indicates that patients with COPD should be screened for OSA due to increased mortality in overlap syndrome, no studies were identified to indicate that CPAP would be prescribed in any manner other than would typically be recommended for patients with clinically significant OSA (see Evidence Review 2.01.18). Patients with overlap syndrome can be treated with CPAP and, when CPAP is not tolerated, with BPAP. The evidence is sufficient to determine that the technology results in an improvement in the net health outcome.

For individuals who have COPD and chronic respiratory failure who receive BPAP, the evidence includes randomized controlled trials (RCTs) and systematic reviews of RCTs. Relevant outcomes are mortality, symptoms, morbid events, functional outcomes, quality of life, and hospitalization. The primary limitation of the evidence base is the heterogeneity of patient selection criteria and treatment parameters. The most recent trials indicate that bilevel non-invasive positive airway pressure ventilation (NPPV) improves hypercapnia in both patients with stable hypercapnia and in patients who have stabilized following an acute exacerbation. There is evidence that some health outcomes including function, readmissions, and death are improved; however, the strength of evidence is low. Several factors have been reported to be important to achieve benefit of NPPV. These are severe hypercapnia with  $\text{PaCO}_2 >52$  mmHg, use for at least 5 hours per night, and treatment with high intensity pressure. In addition, for patients with hypercapnia following an acute exacerbation, titration should occur at least 2 weeks after hospitalization when hypercapnia has stabilized. Under these conditions, the evidence is sufficient to determine that the technology results in an improvement in the net health outcome.

For individuals who have COPD and chronic respiratory failure when BPAP is inadequate who receive home mechanical ventilation (HMV), the evidence includes observational studies and an analysis of administrative claims data. Relevant outcomes are mortality, symptoms, morbid events, functional outcomes, quality of life, and hospitalization. There is low strength of evidence based on observational studies and claims data that NPPV reduces the number of hospital admissions or number of patients with hospitalization compared to either no device or BPAP. Due to the severity of the condition, high quality prospective controlled trials are unlikely in patients who have failed BPAP. HMV may be appropriate in situations where BPAP is not adequate to obtain needed pressures or when daytime use and battery backup is needed. The evidence is sufficient to determine that the technology results in an improvement in the net health outcome.

For individuals who have hypoventilation as a result of thoracic disorder due to neuromuscular disease who receive BPAP, the evidence includes systematic reviews. Relevant outcomes are mortality, symptoms, morbid events, functional outcomes, quality of life, and hospitalization. Clinical trials included in a systematic review of 10 RCTs (or quasi-RCTs) evaluated the use of nocturnal NPPV (primarily BPAP) in individuals with neuromuscular or chest wall disorders. One-year mortality rates were significantly reduced with NPPV use (risk ratio, 0.62; 95% confidence interval [CI], 0.42 to 0.91). Patients treated with NPPV also had lower hospital admission rates and greater symptom improvement. Although the studies were limited by heterogeneity, nocturnal NPPV was found to improve outcomes in patients with restrictive thoracic disorders including neuromuscular disease. A systematic review of observational studies in children with neuromuscular diseases found improved mortality with NPPV compared with standard of care. The evidence is sufficient to determine that the technology results in an improvement in the net health outcome.

For individuals who have hypoventilation as a result of thoracic disorder due to neuromuscular disease who receive HMV, the evidence includes a systematic review of observational studies. One observational study comparing BPAP to HMV found no difference in survival between these ventilation methods, although more patients received effective ventilation with HMV. Due to the severity of the condition, high quality prospective controlled trials are unlikely in patients who have failed BPAP. HMV may be appropriate in situations where BPAP is not adequate to obtain needed pressures or when daytime use and battery backup is needed. The evidence is sufficient to determine that the technology results in an improvement in the net health outcome.

For individuals who have obesity hypoventilation syndrome (OHS) and OSA who receive CPAP, the evidence includes RCTs. In the largest RCT,  $\text{PaCO}_2$  improved with both CPAP and NPPV (mixed BPAP/mechanical ventilation) compared with lifestyle interventions. There was no significant difference between CPAP and NPPV in  $\text{PaCO}_2$  in the short-term or in hospitalized days at long-term follow-up. An RCT comparing CPAP and BPAP found similar outcomes with these treatments. The evidence is sufficient to determine that the technology results in an improvement in the net health outcome.

For individuals who have OHS and chronic respiratory failure who receive BPAP, the evidence includes systematic reviews and RCTs. Relevant outcomes are mortality, symptoms, morbid events, functional outcomes, quality of life, and hospitalization. The majority of evidence specific to BPAP in OHS without OSA comes from a single RCT. In patients with OHS without OSA, BPAP resulted in better PaCO<sub>2</sub> outcomes than lifestyle modifications in the short-term, but long-term outcomes failed to find significant improvement in hospitalization days between groups. The evidence is sufficient to determine that the technology results in an improvement in the net health outcome.

For individuals who have OHS and chronic respiratory failure when BPAP is inadequate who receive HMV, there are no randomized or nonrandomized studies. Due to the severity of the condition, high quality prospective controlled trials are unlikely in patients who have failed BPAP. HMV may be appropriate in situations where BPAP is not adequate to obtain needed pressures or when daytime use and battery backup is needed. The evidence is sufficient to determine that the technology results in an improvement in the net health outcome.

For individuals with hypoventilation as a result of hypoventilation syndromes unrelated to OHS, the evidence is limited to case reports and case series primarily in patients with congenital central hypoventilation. In some cases NPPV minimized the need for invasive mechanical ventilation. Due to the severity of the condition, high quality prospective controlled trials are unlikely in patients who have hypoventilation due to hypoventilation syndromes. The evidence is sufficient to determine that the technology results in an improvement in the net health outcome.

## SUPPLEMENTAL INFORMATION

### Practice Guidelines and Position Statements

Guidelines or position statements will be considered for inclusion in 'Supplemental Information' if they were issued by, or jointly by, a US professional society, an international society with US representation, or National Institute for Health and Care Excellence (NICE). Priority will be given to guidelines that are informed by a systematic review, include strength of evidence ratings, and include a description of management of conflict of interest.

#### American College of Chest Physicians

In 2023, the American College of Chest Physicians (ACCP) published clinical practice guidelines for respiratory management of patients with neuromuscular weakness.<sup>51</sup> Most evidence is based on observational data from patients with amyotrophic lateral sclerosis. The guidelines recommend non-invasive ventilation (NIV) for patients with neuromuscular disease and chronic respiratory failure for patients who meet the following pulmonary function test criteria:

- Forced vital capacity (FVC) <80% predicted with symptoms or FVC <50% predicted without symptoms;
- Maximum inspiratory pressure (MIP) <60 cm H<sub>2</sub>O or maximum expiratory pressure (MEP) <40 cm H<sub>2</sub>O;
- Peak cough flow (PCF) <270 L/min for age ≥12 years or PCF <5th percentile for age <12 years;
- Sniff nasal inspiratory pressure (SNIP) <70 cm H<sub>2</sub>O in male patients, SNIP <60 cmH<sub>2</sub>O in female patients for age ≥12 years.

The panel found no strong evidence to support one method of NIV over another.

#### American College of Chest Physicians et al

In 2021, the ACCP, the American Association for Respiratory Care, the American Academy of Sleep Medicine, and the American Thoracic Society published a technical expert panel report on optimal NIV for chronic obstructive pulmonary disease (COPD), thoracic restrictive disorders, and hypoventilation syndromes.<sup>52,53,54</sup>

##### **Chronic Obstructive Pulmonary Disease**

For COPD the panel recommends that overnight oxygen saturation should not be part of the criteria for bilevel positive airway pressure (BPAP) and that home mechanical ventilators be considered when patients need any of the following:<sup>52</sup>

- "Higher inspiratory pressures than those deliverable by E0471,
- FIO<sub>2</sub> [fraction of inspired oxygen] higher than 40% or 5 L/min nasally,
- Ventilator support for 10 h per day or greater (ie, daytime use),



- Both sophisticated alarms and accompanying internal battery (high-dependency patient),
- Mouthpiece ventilation during the day,
- Persistence of hypercapnia with PaCO<sub>2</sub> [arterial blood carbon dioxide] > 52 mmHg despite adequate adherence to BPAP therapy"
- The panel strongly recommended the use of respiratory therapists in the home for initiation and ongoing support for positive pressure ventilation with either BPAP or home ventilators.

### **Thoracic Restrictive Disorders**

For thoracic restrictive disorders, the panel recommends BPAP for patients with any of the following:<sup>53</sup>,

- "Spirometry (upright or supine) with vital capacity <50% predicted or <80% predicted with associated symptoms (i.e., orthopnea, dyspnea, morning headaches, excessive daytime sleepiness, or unrefreshing sleep),
- Force testing with maximal inspiratory pressure <60 cm H<sub>2</sub>O,
- Hypercapnia:
  - Chronic stable daytime (awake) hypercapnia with PaCO<sub>2</sub> >45 mmHg,
  - Venous blood gas PCO<sub>2</sub>, end-tidal PCO<sub>2</sub>, or transcutaneous PCO<sub>2</sub>, >50 mmHg, or
- Hypoxia:
  - Overnight oximetry in-laboratory or home sleep test with saturation <88% for 5 minutes."

Home mechanical ventilation is recommended in patients with vital capacity <30% or if BPAP fails.

### **Hypoventilation Syndromes**

For patients with hypoventilation syndromes who are obese the recommendations include:<sup>54</sup>,

- BPAP (spontaneous/timed) or volume-assured pressure support (VAPS) for those who are discharged from the hospital, for those with obesity hypoventilation syndrome (OHS) without obstructive sleep apnea, and for those who have failed continuous positive airway pressure (CPAP).

For patients with hypoventilation syndromes due to reduced respiratory drive or advanced lung disease that is not COPD, BPAP (spontaneous/timed) or VAPS is recommended. Patients with hypoventilation syndromes who fail BPAP/VAPS should receive home mechanical ventilation.

## **American Thoracic Society**

### **Chronic Obstructive Pulmonary Disease**

In 2020, the American Thoracic Society published an evidence-based clinical practice guideline on long-term non-invasive ventilation in chronic stable hypercapnic COPD.<sup>2</sup> The society included the recommendations in Table 3, all of which were conditional due to moderate to very low certainty in the evidence base.

**Table 3. American Thoracic Society Recommendations for COPD**

<b>Recommendation</b>	<b>Strength of Recommendation</b>	<b>Level of Certainty</b>
"We suggest the use of nocturnal noninvasive ventilation (NIV) in addition to usual care for patients with chronic stable hypercapnic COPD."	Conditional	Moderate
"We suggest that patients with chronic stable hypercapnic COPD undergo screening for obstructive sleep apnea before initiation of long-term NIV."	Conditional	Very low

"We suggest not initiating long-term NIV during an admission for acute on-chronic hypercapnic respiratory failure, favoring instead reassessment for NIV at 2 - 4 weeks after resolution."	Conditional	Low
"We suggest not using an in-laboratory overnight polysomnogram (PSG) to titrate NIV in patients with chronic stable hypercapnic COPD who are initiating NIV."	Conditional	Very low
"We suggest NIV with targeted normalization of PaCO <sub>2</sub> in patients with hypercapnic COPD on long-term NIV."	Conditional	Low

COPD: chronic obstructive pulmonary disease; NIV: non-invasive ventilation; PaCO<sub>2</sub>: pressure of carbon dioxide; PSG: polysomnogram.  
Hypercapnic COPD defined as PaCO<sub>2</sub> > 45 mmHg.

### **Obesity Hypoventilation Syndrome**

In 2019, the American Thoracic Society published a clinical practice guideline on OHS.<sup>55</sup> These guidelines recommend positive airway pressure for patients with OHS. Generally CPAP is recommended over other NIV because the majority (>70%) of patients have concomitant obstructive sleep apnea (OSA). The guidelines do recommend non-invasive positive airway pressure ventilation (NPPV) initiation at discharge for patients hospitalized with respiratory failure suspected of having OHS until they undergo outpatient workup and titration of positive airway pressure therapy. Both recommendations were conditional with very low level of certainty in the evidence.

### **Global Initiative for Chronic Obstructive Pulmonary Disease**

The Global Initiative for Chronic Obstructive Pulmonary Disease (GOLD) published a revised report for 2024.<sup>1</sup> GOLD guidelines recommend at least 1 of the following as an indication for non-invasive mechanical ventilation :

- Respiratory acidosis (PaCO<sub>2</sub> ≥45 mmHg and arterial pH ≤7.35);
- Severe dyspnea with clinical signs suggestive of respiratory muscle fatigue, increased work of breathing, or both;
- Persistent hypoxemia despite supplemental oxygen therapy.

### **National Institute for Health and Care Excellence Global**

In 2019, the United Kingdom's NICE published a guideline for the diagnosis and management of COPD.<sup>56</sup> NICE recommends that patients with COPD who have chronic hypercapnic respiratory failure despite adequate pharmacologic and oxygen therapy should be referred to a specialist center for consideration of long-term, non-invasive ventilation.

### **U.S. Preventive Services Task Force Recommendations**

Not applicable.

### **Medicare National Coverage**

There is no national coverage determination. In the absence of a national coverage determination, coverage decisions are left to the discretion of local Medicare carriers.

The Centers for Medicare and Medicaid Services requested topic review by the Agency for Healthcare Research and Quality (AHRQ). The technology assessment was published February 2020.<sup>21</sup>

## **REFERENCES**

1. Global Initiative for Chronic Obstructive Lung Disease (GOLD). Global Strategy for Prevention, Diagnosis, and Management of COPD: 2024 Report. [www.goldcopd.org](http://www.goldcopd.org). Accessed January 11, 2024.
2. Macrea M, Oczkowski S, Rochweg B, et al. Long-Term Noninvasive Ventilation in Chronic Stable Hypercapnic Chronic Obstructive Pulmonary Disease. An Official American Thoracic Society Clinical Practice Guideline. *Am J Respir Crit Care Med*. Aug 15 2020; 202(4): e74-e87. PMID

32795139

3. Mathews AM, Wysham NG, Xie J, et al. Hypercapnia in Advanced Chronic Obstructive Pulmonary Disease: A Secondary Analysis of the National Emphysema Treatment Trial. *Chronic Obstr Pulm Dis*. Oct 2020; 7(4): 336-345. PMID 32877962
4. Orr JE, Azofra AS, Tobias LA. Management of Chronic Respiratory Failure in Chronic Obstructive Pulmonary Disease: High-Intensity and Low-Intensity Ventilation. *Sleep Med Clin*. Dec 2020; 15(4): 497-509. PMID 33131660
5. Martinez-Pitre PJ, Sabbula BR, Cascella M. Restrictive Lung Disease. In: *StatPearls*. Treasure Island (FL): StatPearls Publishing; July 25, 2022.
6. Carmona H, Graustein AD, Benditt JO. Chronic Neuromuscular Respiratory Failure and Home Assisted Ventilation. *Annu Rev Med*. Jan 27 2023; 74: 443-455. PMID 36706747
7. Gay PC, Owens RL, Gay PC, et al. Executive Summary: Optimal NIV Medicare Access Promotion: A Technical Expert Panel Report From the American College of Chest Physicians, the American Association for Respiratory Care, the American Academy of Sleep Medicine, and the American Thoracic Society. *Chest*. Nov 2021; 160(5): 1808-1821. PMID 34339685
8. McConville JF, Solway J, Mokhlesi B. Disorders of Ventilation. In: Loscalzo J, Fauci A, Kasper D, Hauser S, Longo D, Jameson J. eds. *Harrison's Principles of Internal Medicine*, 21e. McGraw Hill; 2022. Accessed January 11, 2024. <https://accessmedicine.mhmedical.com/content.aspx?bookid=3095&sectionid=265457022>
9. Wiles SP, Aboussouan LS, Mireles-Cabodevila E. Noninvasive positive pressure ventilation in stable patients with COPD. *Curr Opin Pulm Med*. Mar 2020; 26(2): 175-185. PMID 31895118
10. Srivali N, Thongprayoon C, Tangpanithandee S, et al. The use of continuous positive airway pressure in COPD-O SA overlap syndrome: A systematic review. *Sleep Med*. Aug 2023; 108: 55-60. PMID 37336060
11. Marin JM, Soriano JB, Carrizo SJ, et al. Outcomes in patients with chronic obstructive pulmonary disease and obstructive sleep apnea: the overlap syndrome. *Am J Respir Crit Care Med*. Aug 01 2010; 182(3): 325-31. PMID 20378728
12. Machado MC, Vollmer WM, Togeiro SM, et al. CPAP and survival in moderate-to-severe obstructive sleep apnoea syndrome and hypoxaemic COPD. *Eur Respir J*. Jan 2010; 35(1): 132-7. PMID 19574323
13. Jaoude P, Kufel T, El-Solh AA. Survival benefit of CPAP favors hypercapnic patients with the overlap syndrome. *Lung*. Apr 2014; 192(2): 251-8. PMID 24452812
14. Singh G, Agarwal A, Zhang W, et al. Impact of PAP therapy on hospitalization rates in Medicare beneficiaries with COPD and coexisting OSA. *Sleep Breath*. Mar 2019; 23(1): 193-200. PMID 29931497
15. Raveling T, Vonk J, Struik FM, et al. Chronic non-invasive ventilation for chronic obstructive pulmonary disease. *Cochrane Database Syst Rev*. Aug 09 2021; 8(8): CD002878. PMID 34368950
16. Wilson ME, Dobler CC, Morrow AS, et al. Association of Home Noninvasive Positive Pressure Ventilation With Clinical Outcomes in Chronic Obstructive Pulmonary Disease: A Systematic Review and Meta-analysis. *JAMA*. Feb 04 2020; 323(5): 455-465. PMID 32016309
17. Khnlein T, Windisch W, Khler D, et al. Non-invasive positive pressure ventilation for the treatment of severe stable chronic obstructive pulmonary disease: a prospective, multicentre, randomised, controlled clinical trial. *Lancet Respir Med*. Sep 2014; 2(9): 698-705. PMID 25066329
18. McEvoy RD, Pierce RJ, Hillman D, et al. Nocturnal non-invasive nasal ventilation in stable hypercapnic COPD: a randomised controlled trial. *Thorax*. Jul 2009; 64(7): 561-6. PMID 19213769
19. Murphy PB, Rehal S, Arbane G, et al. Effect of Home Noninvasive Ventilation With Oxygen Therapy vs Oxygen Therapy Alone on Hospital Readmission or Death After an Acute COPD Exacerbation: A Randomized Clinical Trial. *JAMA*. Jun 06 2017; 317(21): 2177-2186. PMID 28528348
20. Struik FM, Sprooten RT, Kerstjens HA, et al. Nocturnal non-invasive ventilation in COPD patients with prolonged hypercapnia after ventilatory support for acute respiratory failure: a randomised, controlled, parallel-group study. *Thorax*. Sep 2014; 69(9): 826-34. PMID 24781217
21. Wilson M, Wang Z, Dobler C, et al. Noninvasive Positive Pressure Ventilation in the Home. Project ID: PULT0717 (Prepared by the Mayo Clinic Evidence-Based Practice Center under Contract No. HHS2902015000131\_HHSA29032004T). Rockville, MD: Agency for Healthcare Research and Quality. March 2019. <https://www.ahrq.gov/sites/default/files/wysiwyg/research/findings/ta/hmv/hmv-ta-fullreport.pdf>. Accessed January 11, 2024.
22. Vasquez MM, McClure LA, Sherrill DL, et al. Positive Airway Pressure Therapies and Hospitalization in Chronic Obstructive Pulmonary Disease. *Am J Med*. Jul 2017; 130(7): 809-818. PMID 28089799
23. Annane D, Orlikowski D, Chevret S. Nocturnal mechanical ventilation for chronic hypoventilation in patients with neuromuscular and chest wall disorders. *Cochrane Database Syst Rev*. Dec 13 2014; 2014(12): CD001941. PMID 25503955
24. Struik FM, Duiverman ML, Meijer PM, et al. Volume-targeted versus pressure-targeted noninvasive ventilation in patients with chest-wall deformity: a pilot study. *Respir Care*. Oct 2011; 56(10): 1522-5. PMID 21513604
25. Jaye J, Chatwin M, Dayer M, et al. Autotitrating versus standard noninvasive ventilation: a randomised crossover trial. *Eur Respir J*. Mar 2009; 33(3): 566-71. PMID 19251798
26. Bourke SC, Tomlinson M, Williams TL, et al. Effects of non-invasive ventilation on survival and quality of life in patients with amyotrophic lateral sclerosis: a randomised controlled trial. *Lancet Neurol*. Feb 2006; 5(2): 140-7. PMID 16426990
27. Tuggey JM, Elliott MW. Randomised crossover study of pressure and volume non-invasive ventilation in chest wall deformity. *Thorax*. Oct 2005; 60(10): 859-64. PMID 16085730
28. Ward S, Chatwin M, Heather S, et al. Randomised controlled trial of non-invasive ventilation (NIV) for nocturnal hypoventilation in neuromuscular and chest wall disease patients with daytime normocapnia. *Thorax*. Dec 2005; 60(12): 1019-24. PMID 16299118
29. Willson GN, Piper AJ, Norman M, et al. Nasal versus full face mask for noninvasive ventilation in chronic respiratory failure. *Eur Respir J*. Apr 2004; 23(4): 605-9. PMID 15083762

30. Laserna E, Barrot E, Beiztegui A, et al. [Non-invasive ventilation in kyphoscoliosis. A comparison of a volumetric ventilator and a BIPAP support pressure device]. *Arch Bronconeumol*. Jan 2003; 39(1): 13-8. PMID 12550014
31. Jackson CE, Rosenfeld J, Moore DH, et al. A preliminary evaluation of a prospective study of pulmonary function studies and symptoms of hypoventilation in ALS/MND patients. *J Neurol Sci*. Oct 15 2001; 191(1-2): 75-8. PMID 11676995
32. Pinto AC, Evangelista T, Carvalho M, et al. Respiratory assistance with a non-invasive ventilator (Bipap) in MND/ALS patients: survival rates in a controlled trial. *J Neurol Sci*. May 1995; 129 Suppl: 19-26. PMID 7595610
33. Raphael JC, Chevret S, Chastang C, et al. Randomised trial of preventive nasal ventilation in Duchenne muscular dystrophy. French Multicentre Cooperative Group on Home Mechanical Ventilation Assistance in Duchenne de Boulogne Muscular Dystrophy. *Lancet*. Jun 25 1994; 343(8913): 1600-4. PMID 7911921
34. AlBalawi MM, Castro-Codesal M, Featherstone R, et al. Outcomes of Long-Term Noninvasive Ventilation Use in Children with Neuromuscular Disease: Systematic Review and Meta-Analysis. *Ann Am Thorac Soc*. Jan 2022; 19(1): 109-119. PMID 34181865
35. Sancho J, Servera E, Morelot-Panzini C, et al. Non-invasive ventilation effectiveness and the effect of ventilatory mode on survival in ALS patients. *Amyotroph Lateral Scler Frontotemporal Degener*. Mar 2014; 15(1-2): 55-61. PMID 24266679
36. Masa JF, Corral J, Alonso ML, et al. Efficacy of Different Treatment Alternatives for Obesity Hypoventilation Syndrome. Pickwick Study. *Am J Respir Crit Care Med*. Jul 01 2015; 192(1): 86-95. PMID 25915102
37. Masa JF, Bentez I, Snchez-Quiroga MÁ, et al. Long-term Noninvasive Ventilation in Obesity Hypoventilation Syndrome Without Severe OSA: The Pickwick Randomized Controlled Trial. *Chest*. Sep 2020; 158(3): 1176-1186. PMID 32343963
38. Masa JF, Corral J, Caballero C, et al. Non-invasive ventilation in obesity hypoventilation syndrome without severe obstructive sleep apnoea. *Thorax*. Oct 2016; 71(10): 899-906. PMID 27406165
39. Masa JF, Mokhlesi B, Bentez I, et al. Long-term clinical effectiveness of continuous positive airway pressure therapy versus non-invasive ventilation therapy in patients with obesity hypoventilation syndrome: a multicentre, open-label, randomised controlled trial. *Lancet*. Apr 27 2019; 393(10182): 1721-1732. PMID 30935737
40. Howard ME, Piper AJ, Stevens B, et al. A randomised controlled trial of CPAP versus non-invasive ventilation for initial treatment of obesity hypoventilation syndrome. *Thorax*. May 2017; 72(5): 437-444. PMID 27852952
41. Arellano-Maric MP, Hamm C, Duiverman ML, et al. Obesity hypoventilation syndrome treated with non-invasive ventilation: Is a switch to CPAP therapy feasible?. *Respirology*. Apr 2020; 25(4): 435-442. PMID 31597227
42. Xu J, Wei Z, Li W, et al. Effect of different modes of positive airway pressure treatment on obesity hypoventilation syndrome: a systematic review and network meta-analysis. *Sleep Med*. Mar 2022; 91: 51-58. PMID 35272117
43. Afshar M, Brozek JL, Soghier I, et al. The Role of Positive Airway Pressure Therapy in Adults with Obesity Hypoventilation Syndrome. A Systematic Review and Meta-Analysis. *Ann Am Thorac Soc*. Mar 2020; 17(3): 344-360. PMID 31726017
44. Borel JC, Tamisier R, Gonzalez-Bermejo J, et al. Noninvasive ventilation in mild obesity hypoventilation syndrome: a randomized controlled trial. *Chest*. Mar 2012; 141(3): 692-702. PMID 21885724
45. Murphy PB, Davidson C, Hind MD, et al. Volume targeted versus pressure support non-invasive ventilation in patients with super obesity and chronic respiratory failure: a randomised controlled trial. *Thorax*. Aug 2012; 67(8): 727-34. PMID 22382596
46. Patout M, Gagnadoux F, Rabec C, et al. AVAPS-AE versus ST mode: A randomized controlled trial in patients with obesity hypoventilation syndrome. *Respirology*. Oct 2020; 25(10): 1073-1081. PMID 32052923
47. Piper AJ, Wang D, Yee BJ, et al. Randomised trial of CPAP vs bilevel support in the treatment of obesity hypoventilation syndrome without severe nocturnal desaturation. *Thorax*. May 2008; 63(5): 395-401. PMID 18203817
48. Storre JH, Seuthe B, Fiechter R, et al. Average volume-assured pressure support in obesity hypoventilation: A randomized crossover trial. *Chest*. Sep 2006; 130(3): 815-21. PMID 16963680
49. Kam K, Bjornson C, Mitchell I. Congenital central hypoventilation syndrome; safety of early transition to non-invasive ventilation. *Pediatr Pulmonol*. Apr 2014; 49(4): 410-3. PMID 23843332
50. Yang L, Qiu S, Zhong J, et al. Noninvasive ventilation via bilevel positive airway pressure improved sleep in a child with congenital central hypoventilation syndrome: A case report. *Clin Case Rep*. Oct 2022; 10(10): e6320. PMID 36276908
51. Khan A, Frazer-Green L, Amin R, et al. Respiratory Management of Patients With Neuromuscular Weakness: An American College of Chest Physicians Clinical Practice Guideline and Expert Panel Report. *Chest*. Aug 2023; 164(2): 394-413. PMID 36921894
52. Hill NS, Criner GJ, Branson RD, et al. Optimal NIV Medicare Access Promotion: Patients With COPD: A Technical Expert Panel Report From the American College of Chest Physicians, the American Association for Respiratory Care, the American Academy of Sleep Medicine, and the American Thoracic Society. *Chest*. Nov 2021; 160(5): e389-e397. PMID 34339684
53. Wolfe LF, Benditt JO, Aboussouan L, et al. Optimal NIV Medicare Access Promotion: Patients With Thoracic Restrictive Disorders: A Technical Expert Panel Report From the American College of Chest Physicians, the American Association for Respiratory Care, the American Academy of Sleep Medicine, and the American Thoracic Society. *Chest*. Nov 2021; 160(5): e399-e408. PMID 34339688
54. Mokhlesi B, Won CH, Make BJ, et al. Optimal NIV Medicare Access Promotion: Patients With Hypoventilation Syndromes: A Technical Expert Panel Report From the American College of Chest Physicians, the American Association for Respiratory Care, the American Academy of Sleep Medicine, and the American Thoracic Society. *Chest*. Nov 2021; 160(5): e377-e387. PMID 34339686
55. Mokhlesi B, Masa JF, Brozek JL, et al. Evaluation and Management of Obesity Hypoventilation Syndrome. An Official American Thoracic Society Clinical Practice Guideline. *Am J Respir Crit Care Med*. Aug 01 2019; 200(3): e6-e24. PMID 31368798
56. National Institute for Health and Care Excellence (NICE). Chronic obstructive pulmonary disease in over 16s: diagnosis and management [NG115]. Updated July 2019 <https://www.nice.org.uk/guidance/ng115>. Accessed January 11, 2024.

## POLICY HISTORY - THIS POLICY WAS APPROVED BY THE FEP® PHARMACY AND MEDICAL POLICY COMMITTEE ACCORDING TO THE HISTORY BELOW:

Date	Action	Description
March 2022	New policy	Policy created with literature review through November 9, 2021. Considered medically necessary under specified conditions.
July 2022	Replace policy	Criteria for second policy statement refined from "after resolution" to "after discharge from the hospital". Clarification added to second Policy Guideline: "Compliance with treatment of at least 4 hours per 24 hours should be documented after the first 3 months of use. There are limited data on which to base compliance assessment. A percentage of days used could be considered, whereby assessment could be based on an average of at least 4 hours per 24 hours over a consecutive 30-day period or use of 4 hours per 24 hours for at least 65% of the days in a consecutive 30-day period." Intent unchanged.
June 2023	Replace policy - correction only	Updated title for related policy 2.01.18.
December 2023	Replace policy	Policy updated with literature review through May 8, 2023; references added. Title changed to remove "Chronic Obstructive Pulmonary Disease." Medically necessary policy statements added for thoracic restrictive disorders due to neuromuscular disease and hypoventilation syndromes for individuals who meet criteria.
June 2024	Replace policy	Policy updated with literature review through January 11, 2024; reference added. Policy statements unchanged.

The policies contained in the FEP Medical Policy Manual are developed to assist in administering contractual benefits and do not constitute medical advice. They are not intended to replace or substitute for the independent medical judgment of a practitioner or other health care professional in the treatment of an individual member. The Blue Cross and Blue Shield Association does not intend by the FEP Medical Policy Manual, or by any particular medical policy, to recommend, advocate, encourage or discourage any particular medical technologies. Medical decisions relative to medical technologies are to be made strictly by members/patients in consultation with their health care providers. The conclusion that a particular service or supply is medically necessary does not constitute a representation or warranty that the Blue Cross and Blue Shield Service Benefit Plan covers (or pays for) this service or supply for a particular member.