

Federal Employee Program® Federal Employee Program® 750 9th St NW Washington, D.C. 20001 202.942.1000 Fax 202.942.1125

5.70.069

Section: Prescription Drugs Effective Date: April 1, 2024

Subsection: Analgesics and Anesthetics Original Policy Date: June 22, 2018

Subject: Olumiant Page: 1 of 10

Last Review Date: March 8, 2024

Olumiant

Description

Olumiant (baricitinib)

Background

Olumiant (baricitinib) is a Janus kinase (JAK) inhibitor. JAKs are intracellular enzymes which transmit signals arising from cytokine or growth factor-receptor interactions on the cellular membrane to influence cellular processes or hematopoiesis and immune cell function. Janus kinase inhibitors inhibit one or more Janus family of enzymes (JAK1, JAK2, JAK3, TYK2), interfering with the JAK-STAT signaling pathway. Within the signaling pathway, JAKs phosphorylate and activate Signal Transducers and Activators of Transcription (STATs) which modulate intracellular activity including gene expression. Olumiant modulates the signaling pathway at the point of JAKs, preventing the phosphorylation and activation of STATs (1).

The use of Olumiant for the treatment of COVID-19 should be billed under the medical benefit.

The use of Olumiant for the treatment of alopecia areata is excluded from coverage.

Regulatory Status

FDA-approved indications: Olumiant is a Janus kinase (JAK) inhibitor indicated for: (1)

- the treatment of adult patients with moderately to severely active rheumatoid arthritis
 (RA) who have had an inadequate response to one or more tumor necrosis factor (TNF)
 blockers.
 - <u>Limitations of Use:</u> Olumiant should not be used in combination with other JAK inhibitors, biological DMARDs or potent immunosuppressants such as azathioprine and cyclosporine.

Section:Prescription DrugsEffective Date:April 1, 2024Subsection:Analgesics and AnestheticsOriginal Policy Date:June 22, 2018

Subject: Olumiant Page: 2 of 10

• the treatment of COVID-19 in hospitalized adult patients requiring supplemental oxygen, non-invasive or invasive mechanical ventilation, or ECMO.

- the treatment of adult patients with severe alopecia areata.
 - <u>Limitations of Use:</u> Olumiant should not be used in combination with other JAK inhibitors, biologic immunomodulators, cyclosporine, or other potent immunosuppressants.

In addition, the FDA has approved the emergency use of Olumiant for the treatment of COVID-19 in hospitalized pediatric patients (2).

Olumiant carries several boxed warnings: (1)

- 1. Serious infections
 - a. Increased risk for serious infections, including tuberculosis and bacterial, invasive fungi, viral and other opportunistic infections that may lead to hospitalization. If a serious infection develops, interrupt Olumiant until the infection is controlled. Prior to the initiation of Olumiant, a test for latent tuberculosis must be conducted. If the test is positive, start treatment for tuberculosis prior to starting Olumiant. Monitor all patients for active tuberculosis during treatment, even if the initial latent tuberculosis test is negative.

2. Mortality

a. RA patients 50 years of age and older with at least one cardiovascular risk factor showed a higher rate of all-cause mortality, including sudden cardiovascular death, in patients treated with JAK inhibitors compared to TNF blockers.

3. Malignancies

- a. Lymphoma and other malignancies have been observed in patients treated with Olumiant.
- b. In RA patients treated with a JAK inhibitor, a higher rate of malignancies was observed when compared with TNF blockers.
- 4. Major adverse cardiovascular events (MACE)
 - a. RA patients 50 years of age and older with at least one cardiovascular risk factor treated with a JAK inhibitor showed a higher rate of MACE (defined as cardiovascular death, myocardial infarction, and stroke) when compared to TNF blockers. Patients who are current or past smokers are at increased risk. Olumiant should be discontinued in patients that have experienced a myocardial infarction or stroke.

5. Thrombosis

Section:Prescription DrugsEffective Date:April 1, 2024Subsection:Analgesics and AnestheticsOriginal Policy Date:June 22, 2018

Subject: Olumiant Page: 3 of 10

a. Thrombosis, including deep vein thrombosis, pulmonary embolism, and arterial thrombosis, have occurred in patients treated with Olumiant. Many of these adverse events were serious and some resulted in death.

b. In RA patients 50 years of age and older with at least one cardiovascular risk factor treated with a JAK inhibitor, a higher rate of thrombosis was observed when compared with TNF blockers.

The safety and effectiveness of Olumiant in pediatric patients have not been established (1).

Related policies

Rinvoq, Xeljanz/XR

Policy

This policy statement applies to clinical review performed for pre-service (Prior Approval, Precertification, Advanced Benefit Determination, etc.) and/or post-service claims.

Olumiant may be considered **medically necessary** if the conditions indicated below are met.

Olumiant may be considered **investigational** for all other indications.

Prior-Approval Requirements

The use of Olumiant for the treatment of COVID-19 should be billed under the medical benefit.

Age 18 years of age or older

Diagnosis

Patient must have the following:

Moderately to severely active rheumatoid arthritis (RA)

AND ALL of the following:

Section:Prescription DrugsEffective Date:April 1, 2024Subsection:Analgesics and AnestheticsOriginal Policy Date:June 22, 2018

Subject: Olumiant Page: 4 of 10

 Inadequate treatment response, intolerance, or contraindication to a 3month trial of at least **ONE** conventional disease-modifying antirheumatic drug (DMARD) (see Appendix 1)

- Inadequate treatment response, intolerance, or contraindication to at least ONE TNF blocker (e.g., Cimzia, Enbrel, Humira, Remicade, Simponi/Simponi Aria)
- 3. Prescriber has considered the risks for malignancy and major adverse cardiovascular events (MACE) (e.g., advanced age, smoking history, cardiovascular risk factors etc.) and determined that Olumiant therapy is appropriate
- 4. Result for latent TB infection is negative OR result was positive for latent TB and patient completed treatment (or is receiving treatment) for latent TB
- 5. Patient **MUST** have tried the preferred product(s) (see Appendix 2) unless the patient has a valid medical exception (e.g., inadequate treatment response, intolerance, contraindication)

AND NONE of the following:

- 1. Active bacterial, invasive fungal, viral, and other opportunistic infections
- 2. Severe hepatic impairment
- 3. A lymphocyte count less than 500 cells/mm3
- 4. An absolute neutrophil count less than 1000 cells/mm3
- A hemoglobin less than 8 g/dL
- 6. History of thrombotic events including deep vein thrombosis (DVT) or pulmonary embolism (PE)
- Used in combination with any other biologic DMARD or targeted synthetic DMARD (see Appendix 1)
- 8. Used in combination with potent immunosuppressants azathioprine or cyclosporine
- 9. Given concurrently with live vaccines

Prior-Approval Renewal Requirements

The use of Olumiant for the treatment of COVID-19 should be billed under the medical benefit.

Age 18 years of age or older

Diagnosis

Section:Prescription DrugsEffective Date:April 1, 2024Subsection:Analgesics and AnestheticsOriginal Policy Date:June 22, 2018

Subject: Olumiant Page: 5 of 10

Patient must have the following:

Rheumatoid arthritis (RA)

AND ALL of the following:

- 1. Condition has improved or stabilized
- Prescriber has considered the risks for malignancy and major adverse cardiovascular events (MACE) (e.g., advanced age, smoking history, cardiovascular risk factors etc.) and determined that continuation of Olumiant therapy is appropriate
- 3. Patient **MUST** have tried the preferred product(s) (see Appendix 2) unless the patient has a valid medical exception (e.g., inadequate treatment response, intolerance, contraindication)

AND NONE of the following:

- 1. Active bacterial, invasive fungal, viral, and other opportunistic infections
- 2. Used in combination with any other biologic DMARD or targeted synthetic DMARD (see Appendix 1)
- 3. Used in combination with potent immunosuppressants azathioprine or cyclosporine
- 4. Development of thrombotic events (including DVTs or PEs)
- 5. Given concurrently with live vaccines

Policy Guidelines

Pre-PA Allowance

None

Prior-Approval Limits

Quantity

Strength	Quantity
1 mg tablet	90 tablets per 90 days
2 mg tablet	90 tablets per 90 days

Section:Prescription DrugsEffective Date:April 1, 2024Subsection:Analgesics and AnestheticsOriginal Policy Date:June 22, 2018

Subject: Olumiant Page: 6 of 10

4 mg tablet	Reserved for the treatment of COVID-19 under the medical benefit	
	OR	
	For the treatment of alopecia areata which is excluded from	
coverage		

Duration 12 months

Prior-Approval Renewal Limits

Quantity

Strength	Quantity	
1 mg tablet	90 tablets per 90 days	
2 mg tablet	90 tablets per 90 days	
4 mg tablet	Reserved for the treatment of COVID-19 under the medical benefit	
	OR	
	For the treatment of alopecia areata which is excluded from	
	coverage	

Duration 18 months

Rationale

Summary

Olumiant (baricitinib) is indicated for the treatment of adult patients with rheumatoid arthritis (RA), for the treatment of COVID-19 in hospitalized patients, and for the treatment of adults with severe alopecia areata. Olumiant has several boxed warnings including increased risk for: serious infections, mortality, malignancies, MACE, and thrombosis. The safety and effectiveness of Olumiant in pediatric patients have not been established (1).

The use of Olumiant for the treatment of COVID-19 should be billed under the medical benefit.

The use of Olumiant for the treatment of alopecia areata is excluded from coverage.

Prior approval is required to ensure the safe, clinically appropriate, and cost-effective use of Olumiant while maintaining optimal therapeutic outcomes.

Section:Prescription DrugsEffective Date:April 1, 2024Subsection:Analgesics and AnestheticsOriginal Policy Date:June 22, 2018

Subject: Olumiant Page: 7 of 10

References

1. Olumiant [package insert]. Indianapolis, IN: Lilly USA, LLC; June 2022.

2. Baricitinib EUA Letter of Authorization. May 2022. Available at: https://www.fda.gov/media/143822/download?utm_medium=email&utm_source=govdelivery

	-	
	Policy History	
	Date	Action
	June 2018	Addition to PA
	September 2018	Annual editorial review
		Addition of requirement of inadequate response, intolerance, or contraindication to a 3-month trial of at least one biological disease-modifying antirheumatic drug (DMARD) per SME
	March 2019	Annual review
	October 2019	Revised quantity limit to 90 tablets per 90 days with the addition of 1 mg tablet availability
	December 2019	Annual review. Addition of requirement to trial preferred product
	March 2020	Annual review and reference update
	December 2020	Annual review and reference update. Added Appendix 2 with a list of
		preferred medications based on diagnosis and plan
	June 2021	Annual review
	January 2022	Added t/f requirement to t/f at least one TNF blocker and added requirement for prescriber to assess risks with malignancy and MACE, per latest PI update
	March 2022	Annual review
	May 2022	COVID-19 added as a new indication per latest PI update. Added statement indicating that the use of Olumiant for COVID-19 must be billed under the medical benefit. Added quantity limit chart statement that the 4 mg tablet is reserved for the treatment of COVID-19 under the medical benefit
	July 2022	Per PI update – addition of alopecia areata to criteria noting that it is excluded from coverage
	September 2022	Annual review
	September 2023	Annual review
	December 2023	Annual review
	March 2024	Annual review
	Keywords	

Section:Prescription DrugsEffective Date:April 1, 2024Subsection:Analgesics and AnestheticsOriginal Policy Date:June 22, 2018

Subject: Olumiant Page: 8 of 10

This policy was approved by the FEP® Pharmacy and Medical Policy Committee on March 8, 2024 and is effective on April 1, 2024.

Section:Prescription DrugsEffective Date:April 1, 2024Subsection:Analgesics and AnestheticsOriginal Policy Date:June 22, 2018

Subject: Olumiant **Page:** 9 of 10

Appendix 1 - List of DMARDS

Conventional disease-modifying antirheumatic drugs (DMARDs)

Generic Name	Brand Name	
azathioprine	Azasan, Imuran	
cyclophosphamide	Cytoxan	
cyclosporine	Neoral, Gengraf, Sandimmune	
hydroxychloroquine	Plaquenil	
leflunomide	Arava	
methotrexate	Rheumatrex, Trexall	
mycophenolate	Cellcept	
sulfasalazine	Azulfidine, Sulfazine	

Biological disease-modifying antirheumatic drugs (DMARDs)

Generic Name	Brand Name	
abatacept	Orencia	
adalimumab	Humira	
anakinra	Kineret	
brodalumab	Siliq	
certolizumab	Cimzia	
etanercept	Enbrel	
golimumab	Simponi/Simponi Aria	
guselkumab	Tremfya	
infliximab	Remicade/Avsola/Inflectra/Renflexis	
ixekizumab	Taltz	
risankizumab-rzaa	Skyrizi	
rituximab	Rituxan/Riabni/Ruxience/Truxima	
sarilumab	Kevzara	
secukinumab	Cosentyx	
spesolimab-sbzo	Spevigo	
tildrakizumab-asmn	Ilumya	
tocilizumab	Actemra	
ustekinumab	Stelara	
vedolizumab	Entyvio	

Targeted synthetic disease-modifying antirheumatic drugs (DMARDs)

Generic Name	Brand Name	
apremilast	Otezla	
baricitinib	Olumiant	
deucravacitinib Sotyktu		

Section:Prescription DrugsEffective Date:April 1, 2024Subsection:Analgesics and AnestheticsOriginal Policy Date:June 22, 2018

Subject: Olumiant Page: 10 of 10

tofacitinib	Xeljanz/XR	
upadactinib	Rinvoq	

Appendix 2 - List of Preferred Products

Diagnosis	Standard Option/Basic Option Preferred Products	Blue Focus Preferred Products
Rheumatoid Arthritis (RA)	*must try TWO preferred products: Actemra SC	*must try ONE preferred product: Enbrel
	Enbrel	Humira**
	Humira** Rinvog	
	Xeljanz/XR	

^{**}Including all preferred biosimilars (see reference product criteria)