

---

# 5.21.156

---

<b>Section:</b>	Prescription Drugs	<b>Effective Date:</b>	April 1, 2026
<b>Subsection:</b>	Antineoplastic Agents	<b>Original Policy Date:</b>	September 4, 2020
<b>Subject:</b>	Monjuvi	<b>Page:</b>	1 of 5

---

**Last Review Date:** March 6, 2026

---

## Monjuvi

### Description

#### Monjuvi (tafasitamab-cxix)

#### Background

Monjuvi (tafasitamab-cxix) is an Fc-modified monoclonal antibody that binds to CD19 antigen expressed on the surface of pre-B and mature B lymphocytes and on several B-cell malignancies, including diffuse large B-cell lymphoma and follicular lymphoma. Upon binding to CD19, Monjuvi mediates B-cell lysis through apoptosis and immune effector mechanisms, including antibody-dependent cellular cytotoxicity and antibody-dependent cellular phagocytosis (1).

#### Regulatory Status

FDA-approved indications: Monjuvi is a CD19-directed cytolytic antibody indicated: (1)

- in combination with lenalidomide for the treatment of adult patients with relapsed or refractory diffuse large B-cell lymphoma (DLBCL) not otherwise specified, including DLBCL arising from low grade lymphoma, and who are not eligible for autologous stem cell transplant (ASCT).
- in combination with lenalidomide and rituximab for the treatment of adult patients with relapsed or refractory follicular lymphoma (FL).

Limitations of use: Monjuvi is not indicated and is not recommended for the treatment of patients with relapsed or refractory marginal zone lymphoma outside of controlled clinical trials (1).

Patients treated for relapsed or refractory DLBCL should receive Monjuvi administered with lenalidomide for a maximum of 12 cycles, then Monjuvi should be continued as monotherapy

# 5.21.156

---

<b>Section:</b>	Prescription Drugs	<b>Effective Date:</b>	April 1, 2026
<b>Subsection:</b>	Antineoplastic Agents	<b>Original Policy Date:</b>	September 4, 2020
<b>Subject:</b>	Monjuvi	<b>Page:</b>	2 of 5

---

until disease progression or unacceptable toxicity. Patients treated for relapsed or refractory FL should be treated with Monjuvi administered in combination with lenalidomide and rituximab for 5 cycles, then Monjuvi should be continued with lenalidomide for a maximum of 7 additional cycles (1).

Monjuvi can cause infusion-related reactions. Patients should be premedicated before starting Monjuvi infusion. Premedications may include acetaminophen, histamine H<sub>1</sub> receptor antagonists, histamine H<sub>2</sub> receptor antagonists, and/or glucocorticosteroids. For patients not experiencing infusion-related reactions during the first 3 infusions, premedication is optional for subsequent infusions (1).

Monjuvi can cause serious or severe myelosuppression, including neutropenia, lymphopenia, thrombocytopenia, and anemia. Complete blood counts (CBC) should be monitored prior to administration of each treatment cycle and throughout treatment (1).

Fatal and serious infections, including opportunistic infections, occurred in patients during treatment with Monjuvi and following the last dose. Patients should be monitored for signs and symptoms of infection and managed as appropriate (1).

Monjuvi may cause fetal B-cell depletion when administered to a pregnant woman. Pregnant women should be advised of the potential risk to a fetus. Females of reproductive potential should be advised to use effective contraception during treatment with Monjuvi and for at least 3 months after the last dose. Monjuvi is initially administered with lenalidomide which is contraindicated in pregnant women because lenalidomide can cause birth defects and death of the unborn child (1).

The safety and effectiveness of Monjuvi in pediatric patients have not been established (1).

## Related policies

Zynlonta

### Policy

*This policy statement applies to clinical review performed for pre-service (Prior Approval, Precertification, Advanced Benefit Determination, etc.) and/or post-service claims.*

Monjuvi may be considered **medically necessary** if the conditions indicated below are met.

Monjuvi may be considered **investigational** for all other indications.

---

<b>Section:</b>	Prescription Drugs	<b>Effective Date:</b>	April 1, 2026
<b>Subsection:</b>	Antineoplastic Agents	<b>Original Policy Date:</b>	September 4, 2020
<b>Subject:</b>	Monjuvi	<b>Page:</b>	3 of 5

---

## Prior-Approval Requirements

**Age** 18 years of age or older

### Diagnoses

Patient must have **ONE** of the following:

1. Relapsed or refractory diffuse large B-cell lymphoma (DLBCL)
  - a. **NOT** eligible for autologous stem cell transplant (ASCT)
  - b. Used in combination with lenalidomide, as tolerated
2. Relapsed or refractory follicular lymphoma (FL)
  - a. Used in combination with lenalidomide and rituximab

**AND ALL** of the following:

- a. Prescriber agrees to monitor the patient for signs and symptoms of infusion-related reactions and administer premedication if needed
- b. Prescriber agrees to monitor complete blood counts (CBC) for myelosuppression
- c. Prescriber agrees to monitor the patient for infections
- d. Female patients of reproductive potential **only**: patient will be advised to use effective contraception during treatment with Monjuvi and for 3 months after the last dose

## Prior – Approval *Renewal* Requirements

**Age** 18 years of age or older

### Diagnosis

Patient must have the following:

1. Relapsed or refractory diffuse large B-cell lymphoma (DLBCL)

**AND ALL** of the following:

- a. **NO** disease progression or unacceptable toxicity
- b. Prescriber agrees to monitor the patient for signs and symptoms of infusion-related reactions and administer premedication if needed

---

<b>Section:</b>	Prescription Drugs	<b>Effective Date:</b>	April 1, 2026
<b>Subsection:</b>	Antineoplastic Agents	<b>Original Policy Date:</b>	September 4, 2020
<b>Subject:</b>	Monjuvi	<b>Page:</b>	4 of 5

---

- c. Prescriber agrees to monitor complete blood counts (CBC) for myelosuppression
- d. Prescriber agrees to monitor the patient for infections
- e. Female patients of reproductive potential **only**: patient will be advised to use effective contraception during treatment with Monjuvi and for 3 months after the last dose

## Policy Guidelines

### Pre – PA Allowance

None

### Prior - Approval Limits

**Duration** 12 months

### Prior – Approval *Renewal* Limits

**Duration** 12 months for DLBCL\*

\***NO** renewal for follicular lymphoma (FL)

## Rationale

### Summary

Monjuvi (tafasitamab-cxix) is an Fc-modified monoclonal antibody that binds to CD19 antigen expressed on the surface of pre-B and mature B lymphocytes and on several B-cell malignancies, including diffuse large B-cell lymphoma and follicular lymphoma. Upon binding to CD19, Monjuvi mediates B-cell lysis through apoptosis and immune effector mechanisms, including antibody-dependent cellular cytotoxicity and antibody-dependent cellular phagocytosis. The safety and effectiveness of Monjuvi in pediatric patients have not been established (1).

Prior approval is required to ensure the safe, clinically appropriate, and cost-effective use of Monjuvi while maintaining optimal therapeutic outcomes.

### References

1. Monjuvi [package insert]. Boston, MA: Morphosys US Inc.; June 2025.

---

<b>Section:</b>	Prescription Drugs	<b>Effective Date:</b>	April 1, 2026
<b>Subsection:</b>	Antineoplastic Agents	<b>Original Policy Date:</b>	September 4, 2020
<b>Subject:</b>	Monjuvi	<b>Page:</b>	5 of 5

---

2. NCCN Drugs & Biologics Compendium<sup>®</sup> Tafasitamab-cxix 2026. National Comprehensive Cancer Network, Inc. Accessed on January 21, 2026.

## Policy History

Date	Action
September 2020	Addition to PA
December 2020	Annual review
June 2021	Annual review and reference update
September 2021	Annual review and reference update
June 2022	Annual review and reference update
March 2023	Annual review and reference update
September 2023	Annual review and reference update
December 2023	Annual review and reference update
March 2024	Annual review and reference update
March 2025	Annual review and reference update
July 2025	Per PI update, added indication of relapsed or refractory follicular lymphoma
September 2025	Annual review and reference update
March 2026	Annual review and reference update

## Keywords

**This policy was approved by the FEP<sup>®</sup> Pharmacy and Medical Policy Committee on March 6, 2026 and is effective on April 1, 2026.**