
5.21.181

Section:	Prescription Drugs	Effective Date:	April 1, 2026
Subsection:	Antineoplastic Agents	Original Policy Date:	September 3, 2021
Subject:	Welireg	Page:	1 of 6

Last Review Date: March 6, 2026

Welireg

Description

Welireg (belzutifan)

Background

Welireg (belzutifan) is an inhibitor of hypoxia-inducible factor 2 alpha (HIF-2 α). HIF-2 α is a transcription factor that plays a role in the body's adaptation response to low oxygen levels. Under normal oxygen levels, HIF-2 α is degraded by the von Hippel-Lindau (VHL) protein. Without functional VHL protein, the HIF-2 α transcription factor accumulates, interacts with hypoxia-inducible factor 1 beta (HIF-1 β) and leads to the expression of genes associated with cellular proliferation, angiogenesis, and tumor growth. Welireg inhibits the formation of the HIF-2 α -HIF-1 β complex, leading to reduced expression of downstream oncogenes (1).

Regulatory Status

FDA-approved indications: Welireg is a hypoxia-inducible factor inhibitor indicated: (1)

- For treatment of adult patients with von Hippel-Lindau (VHL) disease who require therapy for associated renal cell carcinoma (RCC), central nervous system (CNS) hemangioblastomas, or pancreatic neuroendocrine tumors (pNET), not requiring immediate surgery.
- For treatment of adult patients with advanced renal cell carcinoma (RCC) with a clear cell component following a programmed death receptor-1 (PD-1) or programmed death-ligand 1 (PD-L1) inhibitor and a vascular endothelial growth factor tyrosine kinase inhibitor (VEGF-TKI).
- For treatment of adult and pediatric patients 12 years and older with locally advanced, unresectable, or metastatic pheochromocytoma or paraganglioma (PPGL).

5.21.181

Section:	Prescription Drugs	Effective Date:	April 1, 2026
Subsection:	Antineoplastic Agents	Original Policy Date:	September 3, 2021
Subject:	Welireg	Page:	2 of 6

Welireg has a boxed warning regarding embryo-fetal toxicity. Exposure to Welireg during pregnancy can cause embryo-fetal harm and pregnancy status should be verified before initiation of treatment. Welireg can render some hormonal contraceptives ineffective. Female patients of reproductive potential and male patients with partners of reproductive potential should be advised to use effective non-hormonal contraception during treatment with Welireg and for 1 week after the last dose (1).

Welireg has warnings regarding anemia and hypoxia. Patients should be monitored for anemia before initiation and periodically throughout treatment. Welireg should be withheld until hemoglobin $\geq 8\text{g/dL}$, and then resumed at reduced dose or discontinued. Oxygen saturation should be monitored before initiating treatment and then periodically throughout treatment. If patient becomes hypoxic at rest, withhold Welireg until resolved, and then resume at reduced dose or discontinue permanently. In cases of life-threatening hypoxia, discontinue Welireg permanently (1).

The safety and effectiveness of Welireg in pediatric patients less than 12 years of age for the treatment of locally advanced, unresectable, or metastatic pheochromocytoma or paraganglioma have not been established. The safety and effectiveness of Welireg in pediatric patients less than 18 years of age for all other indications have not been established (1).

Related policies

Policy

This policy statement applies to clinical review performed for pre-service (Prior Approval, Precertification, Advanced Benefit Determination, etc.) and/or post-service claims.

Welireg may be considered **medically necessary** if the conditions indicated below are met.

Welireg may be considered **investigational** for all other indications.

Prior-Approval Requirements

Diagnoses

Patient must have **ONE** of the following:

5.21.181

Section:	Prescription Drugs	Effective Date:	April 1, 2026
Subsection:	Antineoplastic Agents	Original Policy Date:	September 3, 2021
Subject:	Welireg	Page:	3 of 6

1. Von Hippel-Lindau (VHL) disease
 - a. 18 years of age or older
 - b. Patient has **ONE** of the following:
 - i. Renal cell carcinoma (RCC)
 - ii. Central nervous system (CNS) hemangioblastomas
 - iii. Pancreatic neuroendocrine tumors (pNET)
 - c. Patient does not require immediate surgery
2. Advanced renal cell carcinoma (RCC)
 - a. 18 years of age or older
 - b. Previous treatment with **ALL** of the following:
 - i. PD-1 inhibitor **OR** PD-L1 inhibitor
 - ii. VEGF-TKI
3. Locally advanced, unresectable, or metastatic pheochromocytoma or paraganglioma (PPGL)
 - a. 12 years of age or older

AND ALL of the following:

1. Hemoglobin ≥ 8 g/dL
2. Prescriber agrees to monitor for anemia and hypoxia before initiation of treatment and periodically throughout treatment
3. Females of reproductive potential **only**: patient has had a negative pregnancy test **AND** patient will be advised to use effective non-hormonal contraception during treatment with Welireg and for 1 week after the last dose
4. Males with female partners of reproductive potential **only**: pregnancy will be excluded before start of treatment and patient will be advised to use effective non-hormonal contraception during treatment with Welireg and for 1 week after the last dose

Prior-Approval *Renewal* Requirements

Diagnoses

Patient must have **ONE** of the following:

1. Von Hippel-Lindau (VHL) disease
 - a. 18 years of age or older

5.21.181

Section:	Prescription Drugs	Effective Date:	April 1, 2026
Subsection:	Antineoplastic Agents	Original Policy Date:	September 3, 2021
Subject:	Welireg	Page:	4 of 6

- b. Patient has **ONE** of the following:
 - i. Renal cell carcinoma (RCC)
 - ii. Central nervous system (CNS) hemangioblastomas
 - iii. Pancreatic neuroendocrine tumors (pNET)
- c. Patient does not require immediate surgery
- 2. Advanced renal cell carcinoma (RCC)
 - a. 18 years of age or older
- 3. Advanced, unresectable, or metastatic pheochromocytoma or paraganglioma (PPGL)
 - a. 12 years of age or older

AND ALL of the following:

- 1. **NO** disease progression or unacceptable toxicity
- 2. Hemoglobin ≥ 8 g/dL
- 3. Prescriber agrees to monitor for anemia and hypoxia periodically throughout treatment
- 4. Females of reproductive potential **only**: patient will be advised to use effective non-hormonal contraception during treatment with Welireg and for 1 week after the last dose
- 5. Males with female partners of reproductive potential **only**: patient will be advised to use effective non-hormonal contraception during treatment with Welireg and for 1 week after the last dose

Policy Guidelines

Pre-PA Allowance

None

Prior-Approval Limits

Quantity 120 mg per day

Duration 12 months

Prior-Approval *Renewal* Limits

Same as above

5.21.181

Section:	Prescription Drugs	Effective Date:	April 1, 2026
Subsection:	Antineoplastic Agents	Original Policy Date:	September 3, 2021
Subject:	Welireg	Page:	5 of 6

Rationale

Summary

Welireg (belzutifan) is an inhibitor of hypoxia-inducible factor 2 alpha (HIF-2 α) and is indicated for von Hippel-Lindau (VHL) disease, advanced renal cell carcinoma (RCC), and locally advanced, unresectable, or metastatic pheochromocytoma or paraganglioma (PPGL). Welireg carries a boxed warning regarding embryo-fetal toxicity and patients should be advised to use to effective non-hormonal contraception. Welireg has also been shown to cause hypoxemia and anemia. Patients should be monitored for these conditions and dosage adjusted, or treatment discontinued as appropriate. The safety and effectiveness of Welireg in pediatric patients less than 12 years of age for the treatment of locally advanced, unresectable, or metastatic pheochromocytoma or paraganglioma have not been established. The safety and effectiveness of Welireg in pediatric patients less than 18 years of age for all other indications have not been established (1).

Prior approval is required to ensure the safe, clinically appropriate, and cost-effective use of Welireg while maintaining optimal therapeutic outcomes.

References

1. Welireg [package insert]. Whitehouse Station, NJ: Merck Sharpe & Dohme Corp.; May 2025.
2. NCCN Drugs & Biologics Compendium[®] Belzutifan 2026. National Comprehensive Cancer Network, Inc. Accessed on January 21, 2026.

Policy History

Date	Action
September 2021	Addition to PA
December 2021	Annual review and reference update
December 2022	Annual review and reference update
March 2023	Annual review and reference update
January 2024	Per PI update, added indication of advanced renal cell carcinoma (RCC). Changed hemoglobin requirement to ≥ 8 g/dL. Changed quantity limit to 120 mg per day
March 2024	Annual review and reference update

5.21.181

Section:	Prescription Drugs	Effective Date:	April 1, 2026
Subsection:	Antineoplastic Agents	Original Policy Date:	September 3, 2021
Subject:	Welireg	Page:	6 of 6

March 2025	Annual editorial review and reference update. Reworded initiation requirement to “negative pregnancy test”
June 2025	Per PI update, added indication of locally advance, unresectable, or metastatic pheochromocytoma or paraganglioma (PPGL)
September 2025	Annual review and reference update
March 2026	Annual review and reference update

Keywords

This policy was approved by the FEP® Pharmacy and Medical Policy Committee on March 6, 2026 and is effective on April 1, 2026.