
5.90.004

| | | | |
|--------------------|--------------------|------------------------------|-------------------|
| Section: | Prescription Drugs | Effective Date: | April 1, 2026 |
| Subsection: | Biologicals | Original Policy Date: | November 15, 2013 |
| Subject: | Stelara | Page: | 1 of 17 |

Last Review Date: March 6, 2026

Stelara

Description

Stelara (ustekinumab)
Imuldosa (ustekinumab-srlf)
Otulfi (**ustekinumab-aauz**)
Pyzchiva (ustekinumab-ttwe)
Selarsdi (ustekinumab-aekn)
Starjemza (ustekinumab-hmny)
Steqeyma (ustekinumab-stba)
Wezlana (ustekinumab-auub)
Yesintek (ustekinumab-kfce)

Bolded medications are the preferred products.

Background

Stelara and its biosimilars are human interleukin-12 (IL-12) and interleukin-23 (IL-23) antagonists indicated for the treatment of plaque psoriasis, psoriatic arthritis, Crohn's disease, and ulcerative colitis. Stelara and its biosimilars targets IL-12 and IL-23, reducing inflammation and relieving symptoms of joint pain, swelling, stiffness, plaque thickness, scaling, and redness

| | | | |
|--------------------|--------------------|------------------------------|-------------------|
| Section: | Prescription Drugs | Effective Date: | April 1, 2026 |
| Subsection: | Biologicals | Original Policy Date: | November 15, 2013 |
| Subject: | Stelara | Page: | 2 of 17 |

in psoriatic arthritis and plaque psoriasis, and has been shown to significantly decrease disease activity in patients with moderately to severely active Crohn's disease and ulcerative colitis (1-9).

Regulatory Status

FDA-approved indications: Stelara and its biosimilars are human interleukin-12 and -23 antagonists indicated for the treatment of: (1-9)

Adult patients with:

1. Moderate to severe plaque psoriasis (PsO) who are candidates for phototherapy or systemic therapy
2. Active psoriatic arthritis (PsA)
3. Moderately to severely active Crohn's disease (CD)
4. Moderately to severely active ulcerative colitis (UC)

Pediatric patients 6 years and older with:

1. Moderate to severe plaque psoriasis, who are candidates for phototherapy or systemic therapy
2. Active psoriatic arthritis (PsA)

Stelara and its biosimilars may increase the risk of infections and reactivation of latent infections such as bacterial, fungal, and viral infections. Stelara and its biosimilars should not be given to patients with any clinically important active infection until the infection resolves or is adequately treated. Serious infections that require hospitalization may occur such as diverticulitis, cellulitis, pneumonia, appendicitis, sepsis, and cholecystitis (1-9).

Evaluate patients for tuberculosis infection prior to initiating treatment with Stelara or its biosimilars. Do not administer Stelara or its biosimilars to patients with active tuberculosis. Initiate treatment of latent tuberculosis prior to administering Stelara or its biosimilars. Consider anti-tuberculosis therapy prior to initiation of Stelara or its biosimilars in patients with a past history of latent or active tuberculosis in whom an adequate course of treatment cannot be confirmed. Patients receiving Stelara or its biosimilars should be monitored closely for signs and symptoms of active tuberculosis during and after treatment (1-9).

Stelara and its biosimilars are immunosuppressants and may increase the risk of malignancy. Malignancies were reported among subjects who received Stelara or its biosimilars. There have been post-marketing reports of the rapid appearance of multiple cutaneous squamous cell carcinomas in patients receiving Stelara or its biosimilars who had pre-existing risk factors for developing non-melanoma skin cancer. All patients receiving Stelara or its biosimilars should be monitored for the appearance of non-melanoma skin cancer. Patients greater than 60 years of

| | | | |
|--------------------|--------------------|------------------------------|-------------------|
| Section: | Prescription Drugs | Effective Date: | April 1, 2026 |
| Subsection: | Biologicals | Original Policy Date: | November 15, 2013 |
| Subject: | Stelara | Page: | 3 of 17 |

age, those with a medical history of prolonged immunosuppressant therapy, and those with a history of PUVA treatment should be followed closely (1-9).

Safety and effectiveness of Stelara and its biosimilars in pediatric patients less than 6 years of age with plaque psoriasis have not been established (1-9).

Safety and effectiveness of Stelara and its biosimilars in pediatric patients less than 18 years of age with psoriatic arthritis, Crohn's disease, or ulcerative colitis have not been established (1-9).

Related policies

Ilumya, Skyrizi, Tremfya

Policy

This policy statement applies to clinical review performed for pre-service (Prior Approval, Precertification, Advanced Benefit Determination, etc.) and/or post-service claims.

Stelara and its biosimilars may be considered **medically necessary** if the conditions indicated below are met.

Stelara and its biosimilars may be considered **investigational** for all other indications.

Prior-Approval Requirements

Preferred medications only

Diagnoses

Patient must have **ONE** of the following:

1. Moderate to severe plaque psoriasis (PsO)
 - a. 6 years of age or older
 - b. Inadequate treatment response, intolerance, or contraindication to either conventional systemic therapy (see Appendix 1) or phototherapy
 - i. If the patient is intolerant or contraindicated to one therapy then the patient must have an inadequate treatment response, intolerance, or contraindication to the other treatment option

| | | | |
|--------------------|--------------------|------------------------------|-------------------|
| Section: | Prescription Drugs | Effective Date: | April 1, 2026 |
| Subsection: | Biologicals | Original Policy Date: | November 15, 2013 |
| Subject: | Stelara | Page: | 4 of 17 |

- c. Prescriber will not exceed the FDA labeled maintenance dose of **ONE** of the following:
 - i. Subcutaneous administration: Patients 6-17 years of age and less than 60 kg weight – 0.75 mg/kg every 12 weeks
 - ii. Subcutaneous administration: Patients 6-17 years of age 60 kg to 100 kg weight and adult patients less than or equal to 100 kg weight – 45 mg every 12 weeks
 - iii. Subcutaneous administration: Patients greater than 100 kg weight – 90 mg every 12 weeks
2. Active psoriatic arthritis (PsA)
 - a. 6 years of age or older
 - b. Inadequate treatment response, intolerance, or contraindication to a 3-month trial of at least **ONE** conventional DMARD (see Appendix 1)
 - c. Prescriber will not exceed the FDA labeled maintenance dose of **ONE** of the following:
 - i. Subcutaneous administration: Patients 18 years of age or older - 45 mg every 12 weeks
 - ii. Subcutaneous administration: Patients 6 years of age or older, weight greater than 100 kg, with concurrent moderate to severe plaque psoriasis – 90 mg every 12 weeks
 - iii. Subcutaneous administration: Patients 6-17 years of age and less than 60 kg weight - 0.75 mg/kg every 12 weeks
 - iv. Subcutaneous administration: Patients 6-17 years of age and greater than or equal to 60 kg weight - 45 mg every 12 weeks
3. Moderate to severely active Crohn's disease (CD)
 - a. 18 years of age or older
 - b. Inadequate treatment response, intolerance, or contraindication to at least **ONE** conventional therapy option (see Appendix 2)
 - c. Prescriber will initiate dosing with a single intravenous infusion with **ONE** of the following:
 - i. IV infusion: 55 kg or less – 260 mg
 - ii. IV infusion: >55 kg to 85 kg – 390 mg
 - iii. IV infusion: More than 85 kg – 520 mg
 - d. Prescriber will not exceed the FDA labeled maintenance dose of the following:
 - i. Subcutaneous administration: 90 mg every 8 weeks

| | | | |
|--------------------|--------------------|------------------------------|-------------------|
| Section: | Prescription Drugs | Effective Date: | April 1, 2026 |
| Subsection: | Biologicals | Original Policy Date: | November 15, 2013 |
| Subject: | Stelara | Page: | 5 of 17 |

4. Moderate to severely active ulcerative colitis (UC)
 - a. 18 years of age or older
 - b. Inadequate treatment response, intolerance, or contraindication to at least **ONE** conventional therapy option (see Appendix 2)
 - c. Prescriber will initiate dosing with a single intravenous infusion with **ONE** of the following:
 - i. IV infusion: 55 kg or less – 260 mg
 - ii. IV infusion: >55 kg to 85 kg – 390 mg
 - iii. IV infusion: More than 85 kg – 520 mg
 - d. Prescriber will not exceed the FDA labeled maintenance dose of the following:
 - i. Subcutaneous administration: 90 mg every 8 weeks

AND ALL of the following for **ALL** diagnoses:

1. Result for latent TB infection is negative **OR** result was positive for latent TB and patient completed treatment (or is receiving treatment) for latent TB
2. Absence of active infection [including tuberculosis and hepatitis B virus (HBV)]
3. **NOT** to be used in combination with any other biologic DMARD or targeted synthetic DMARD (see Appendix 1)
4. **NOT** given concurrently with live vaccines

Non-preferred medications only

Diagnoses

Patient must have **ONE** of the following with provided documentation (e.g., medical records, laboratory reports):

1. Moderate to severe plaque psoriasis (PsO)
 - a. 6 years of age or older
 - b. Inadequate treatment response, intolerance, or contraindication to either conventional systemic therapy (see Appendix 1) or phototherapy
 - i. If the patient is intolerant or contraindicated to one therapy then the patient must have an inadequate treatment response, intolerance, or contraindication to the other treatment option
 - c. Prescriber will not exceed the FDA labeled maintenance dose of **ONE** of the following:

| | | | |
|--------------------|--------------------|------------------------------|-------------------|
| Section: | Prescription Drugs | Effective Date: | April 1, 2026 |
| Subsection: | Biologicals | Original Policy Date: | November 15, 2013 |
| Subject: | Stelara | Page: | 6 of 17 |

- i. Subcutaneous administration: Patients 6-17 years of age and less than 60 kg weight – 0.75 mg/kg every 12 weeks
 - ii. Subcutaneous administration: Patients 6-17 years of age 60 kg to 100 kg weight and adult patients less than or equal to 100 kg weight – 45 mg every 12 weeks
 - iii. Subcutaneous administration: Patients greater than 100 kg weight – 90 mg every 12 weeks
2. Active psoriatic arthritis (PsA)
 - a. 6 years of age or older
 - b. Inadequate treatment response, intolerance, or contraindication to a 3-month trial of at least **ONE** conventional DMARD (see Appendix 1)
 - c. Prescriber will not exceed the FDA labeled maintenance dose of **ONE** of the following:
 - i. Subcutaneous administration: Patients 18 years of age or older - 45 mg every 12 weeks
 - ii. Subcutaneous administration: Patients 6 years of age or older, weight greater than 100 kg, with concurrent moderate to severe plaque psoriasis – 90 mg every 12 weeks
 - iii. Subcutaneous administration: Patients 6-17 years of age and less than 60 kg weight - 0.75 mg/kg every 12 weeks
 - iv. Subcutaneous administration: Patients 6-17 years of age and greater than or equal to 60 kg weight - 45 mg every 12 weeks
3. Moderate to severely active Crohn's disease (CD)
 - a. 18 years of age or older
 - b. Inadequate treatment response, intolerance, or contraindication to at least **ONE** conventional therapy option (see Appendix 2)
 - c. Prescriber will initiate dosing with a single intravenous infusion with **ONE** of the following:
 - i. IV infusion: 55 kg or less – 260 mg
 - ii. IV infusion: >55 kg to 85 kg – 390 mg
 - iii. IV infusion: More than 85 kg – 520 mg
 - d. Prescriber will not exceed the FDA labeled maintenance dose of the following:
 - i. Subcutaneous administration: 90 mg every 8 weeks
4. Moderate to severely active ulcerative colitis (UC)
 - a. 18 years of age or older

| | | | |
|--------------------|--------------------|------------------------------|-------------------|
| Section: | Prescription Drugs | Effective Date: | April 1, 2026 |
| Subsection: | Biologicals | Original Policy Date: | November 15, 2013 |
| Subject: | Stelara | Page: | 7 of 17 |

- b. Inadequate treatment response, intolerance, or contraindication to at least **ONE** conventional therapy option (see Appendix 2)
- c. Prescriber will initiate dosing with a single intravenous infusion with **ONE** of the following:
 - i. IV infusion: 55 kg or less – 260 mg
 - ii. IV infusion: >55 kg to 85 kg – 390 mg
 - iii. IV infusion: More than 85 kg – 520 mg
- d. Prescriber will not exceed the FDA labeled maintenance dose of the following:
 - i. Subcutaneous administration: 90 mg every 8 weeks

AND ALL of the following for **ALL** diagnoses:

- 1. Result for latent TB infection is negative **OR** result was positive for latent TB and patient completed treatment (or is receiving treatment) for latent TB
- 2. Absence of active infection [including tuberculosis and hepatitis B virus (HBV)]
- 3. **NOT** to be used in combination with any other biologic DMARD or targeted synthetic DMARD (see Appendix 1)
- 4. **NOT** given concurrently with live vaccines
- 5. Patient **MUST** have tried the preferred product(s) (see Appendix 3) unless the patient has a valid medical exception (e.g., inadequate treatment response, intolerance, contraindication)

All approved requests are subject to review by a clinical specialist for final validation and coverage determination once all required documentation has been received. Current utilization, including samples, does not guarantee approval of coverage.

Prior – Approval *Renewal* Requirements

Preferred medications only

Diagnoses

Patient must have **ONE** of the following:

- 1. Plaque psoriasis (PsO)
 - a. 6 years of age or older
 - b. Prescriber will not exceed the FDA labeled maintenance dose of **ONE** of the following:

| | | | |
|--------------------|--------------------|------------------------------|-------------------|
| Section: | Prescription Drugs | Effective Date: | April 1, 2026 |
| Subsection: | Biologicals | Original Policy Date: | November 15, 2013 |
| Subject: | Stelara | Page: | 8 of 17 |

- i. Subcutaneous administration: Patients 6-17 years of age and less than 60 kg weight – 0.75 mg/kg every 12 weeks
 - ii. Subcutaneous administration: Patients 6-17 years of age and 60 kg to 100 kg weight and adult patients less than or equal to 100 kg weight – 45 mg every 12 weeks
 - iii. Subcutaneous administration: Patients greater than 100 kg weight – 90 mg every 12 weeks
 2. Psoriatic arthritis (PsA)
 - a. 6 years of age or older
 - b. Prescriber will not exceed the FDA labeled maintenance dose of **ONE** of the following:
 - i. Subcutaneous administration: Patients 18 years of age or older - 45 mg every 12 weeks
 - ii. Subcutaneous administration: Patients 6 years of age or older, weight greater than 100 kg, with concurrent moderate to severe plaque psoriasis – 90 mg every 12 weeks
 - iii. Subcutaneous administration: Patients 6-17 years of age and less than 60 kg weight - 0.75 mg/kg every 12 weeks
 - iv. Subcutaneous administration: Patients 6-17 years of age and greater than or equal to 60 kg weight - 45 mg every 12 weeks
 3. Crohn's disease (CD)
 - a. 18 years of age or older
 - b. Prescriber will not exceed the FDA labeled maintenance dose of the following:
 - i. Subcutaneous administration: 90 mg every 8 weeks
 4. Ulcerative colitis (UC)
 - a. 18 years of age or older
 - b. Prescriber will not exceed the FDA labeled maintenance dose of the following:
 - i. Subcutaneous administration: 90 mg every 8 weeks

AND ALL of the following for **ALL** diagnoses:

1. Condition has improved or stabilized with Stelara
2. Absence of active infection [including tuberculosis and hepatitis B virus (HBV)]
3. **NOT** to be used in combination with any other biologic DMARD or targeted synthetic DMARD (see Appendix 1)

| | | | |
|--------------------|--------------------|------------------------------|-------------------|
| Section: | Prescription Drugs | Effective Date: | April 1, 2026 |
| Subsection: | Biologicals | Original Policy Date: | November 15, 2013 |
| Subject: | Stelara | Page: | 9 of 17 |

4. **NOT** given concurrently with live vaccines
-

Non-preferred medications only

Diagnoses

Patient must have **ONE** of the following with provided documentation (e.g., medical records, laboratory reports):

1. Plaque psoriasis (PsO)
 - a. 6 years of age or older
 - b. Prescriber will not exceed the FDA labeled maintenance dose of **ONE** of the following:
 - i. Subcutaneous administration: Patients 6-17 years of age and less than 60 kg weight – 0.75 mg/kg every 12 weeks
 - ii. Subcutaneous administration: Patients 6-17 years of age and 60 kg to 100 kg weight and adult patients less than or equal to 100 kg weight – 45 mg every 12 weeks
 - iii. Subcutaneous administration: Patients greater than 100 kg weight – 90 mg every 12 weeks
2. Psoriatic arthritis (PsA)
 - a. 6 years of age or older
 - b. Prescriber will not exceed the FDA labeled maintenance dose of **ONE** of the following:
 - i. Subcutaneous administration: Patients 18 years of age or older - 45 mg every 12 weeks
 - ii. Subcutaneous administration: Patients 6 years of age or older, weight greater than 100 kg, with concurrent moderate to severe plaque psoriasis – 90 mg every 12 weeks
 - iii. Subcutaneous administration: Patients 6-17 years of age and less than 60 kg weight - 0.75 mg/kg every 12 weeks
 - iv. Subcutaneous administration: Patients 6-17 years of age and greater than or equal to 60 kg weight - 45 mg every 12 weeks
3. Crohn's disease (CD)
 - a. 18 years of age or older

| | |
|------------------------------------|--|
| Section: Prescription Drugs | Effective Date: April 1, 2026 |
| Subsection: Biologicals | Original Policy Date: November 15, 2013 |
| Subject: Stelara | Page: 10 of 17 |

- b. Prescriber will not exceed the FDA labeled maintenance dose of the following:
 - i. Subcutaneous administration: 90 mg every 8 weeks

- 4. Ulcerative colitis (UC)
 - a. 18 years of age or older
 - b. Prescriber will not exceed the FDA labeled maintenance dose of the following:
 - i. Subcutaneous administration: 90 mg every 8 weeks

AND ALL of the following for **ALL** diagnoses:

1. Condition has improved or stabilized with Stelara
2. Absence of active infection [including tuberculosis and hepatitis B virus (HBV)]
3. **NOT** to be used in combination with any other biologic DMARD or targeted synthetic DMARD (see Appendix 1)
4. **NOT** given concurrently with live vaccines
5. Patient **MUST** have tried the preferred product(s) (see Appendix 3) unless the patient has a valid medical exception (e.g., inadequate treatment response, intolerance, contraindication)

All approved requests are subject to review by a clinical specialist for final validation and coverage determination once all required documentation has been received. Current utilization, including samples, does not guarantee approval of coverage.

Policy Guidelines

Pre - PA Allowance

None

Prior - Approval Limits

Quantity

| Diagnosis | Strength | Quantity |
|---------------------------|--|---|
| Plaque psoriasis (PsO) | <u>Weight ≤100kg</u> 45 mg SC vial/syringe <u>Weight > 100kg</u> 90 mg SC syringe | 5 units per 365 days (dosed initially, 4 weeks later, then every 12 weeks) |
| Psoriatic arthritis (PsA) | 45 mg SC vial/syringe <u>Concurrent moderate to severe plaque psoriasis and weight > 100kg</u> | |

| | | | |
|--------------------|--------------------|------------------------------|-------------------|
| Section: | Prescription Drugs | Effective Date: | April 1, 2026 |
| Subsection: | Biologicals | Original Policy Date: | November 15, 2013 |
| Subject: | Stelara | Page: | 11 of 17 |

| | | |
|-------------------------|------------------------------------|---|
| | 90 mg SC syringe | |
| Crohn's disease (CD) | | <u>Weight ≤55kg</u> 2 IV vials (1 dose) + 1 SC syringe per 56 days |
| Ulcerative colitis (UC) | 130 mg IV vial 90 mg SC syringe | <u>Weight > 55kg to 85kg</u> 3 IV vials (1 dose) + 1 SC syringe per 56 days |
| | | <u>Weight > 85kg</u> 4 IV vials (1 dose) + 1 SC syringe per 56 days |

Duration 12 months

Prior – Approval *Renewal* Limits

Quantity

| Diagnosis | Strength | Quantity |
|---------------------------|--|--------------------------|
| Plaque psoriasis (PsO) | <u>Weight ≤100kg</u> 45 mg SC vial/syringe | 1 unit per 84 days |
| | <u>Weight > 100kg</u> 90 mg SC syringe | |
| Psoriatic arthritis (PsA) | 45 mg SC vial/syringe | 1 SC syringe per 56 days |
| | <u>Concurrent moderate to severe plaque psoriasis and weight > 100kg</u> 90 mg SC syringe | |
| Crohn's disease (CD) | 90 mg SC syringe | |
| Ulcerative colitis (UC) | | |

Duration 18 months

Rationale

Summary

Stelara and its biosimilars are human interleukin-12 (IL-12) and interleukin-23 (IL-23) antagonists indicated for the treatment of plaque psoriasis, psoriatic arthritis, Crohn's disease, and ulcerative colitis. Stelara and its biosimilars target IL-12 and IL-23, reducing inflammation and relieving symptoms of joint pain, swelling, stiffness, plaque thickness, scaling, and redness

| | | | |
|--------------------|--------------------|------------------------------|-------------------|
| Section: | Prescription Drugs | Effective Date: | April 1, 2026 |
| Subsection: | Biologicals | Original Policy Date: | November 15, 2013 |
| Subject: | Stelara | Page: | 12 of 17 |

in psoriatic arthritis and plaque psoriasis, and have been shown to significantly decrease disease activity in patients with moderately to severely active Crohn's disease and ulcerative colitis. Stelara and its biosimilars may increase the risk of infections and reactivation of latent infections such as bacterial, fungal, and viral infections. Stelara and its biosimilars should not be given to patients with any clinically important active infection until the infection resolves or is adequately treated. Stelara and its biosimilars should not be administered to patients with active TB. Stelara and its biosimilars may increase the risk of malignancy (1-9).

Prior authorization is required to ensure the safe, clinically appropriate, and cost-effective use of Stelara and its biosimilar while maintaining optimal therapeutic outcomes.

References

1. Stelara [package insert]. Horsham, PA: Janssen Biotech, Inc.; November 2025.
2. Selarsdi [package insert]. Parsippany, NJ: Teva Pharmaceuticals; December 2025.
3. Pyzchiva [package insert]. Princeton, NJ: Sandoz Inc.; November 2025.
4. Yesintek [package insert]. Cambridge, MA: Biocon Biologics Inc.; November 2024.
5. Steqeyma [package insert]. Jersey City, NJ: Celltrion; December 2025.
6. Wezlana [package insert]. Thousand Oaks, CA: Amgen Inc.; September 2025.
7. Imuldosa [package insert]. Raleigh, NC: Accord BioPharma Inc.; November 2025.
8. Otulfi [package insert]. Lake Zurich, IL: Fresenius Kabi USA, LLC; August 2025.
9. Starjemza [package insert]. Berkeley Heights, NJ: Hikma Pharmaceuticals USA Inc.; May 2025.

Policy History

| Date | Action |
|----------------|--|
| October 2013 | Addition to PA |
| December 2013 | Annual editorial review by the PMPC |
| September 2014 | Annual editorial review and renewal limit to 18 months |
| September 2016 | Annual editorial review and reference update Addition of not to be used in combination with any other biologic DMARD or targeted synthetic DMARD Addition of not given concurrently with live vaccines per SME Policy number change from 5.18.04 to 5.90.04 |
| October 2016 | Addition of Crohn's disease to diagnoses in initiation and renewal criteria Addition of criteria to Crohn's disease diagnosis in initiation: must have inadequate treatment response to one of the following: immunomodulators, corticosteroids, or TNF blockers |
| December 2016 | Annual review |

| | | | |
|--------------------|--------------------|------------------------------|-------------------|
| Section: | Prescription Drugs | Effective Date: | April 1, 2026 |
| Subsection: | Biologicals | Original Policy Date: | November 15, 2013 |
| Subject: | Stelara | Page: | 13 of 17 |

| | |
|----------------|---|
| September 2017 | Annual editorial review |
| | Addition of FDA dosing requirement questions for all indications |
| October 2017 | Addition of PsO dosing for 12 yrs. of age and older |
| December 2017 | Annual review |
| June 2018 | Addition of IV initiation dosing for CD |
| | Addition of additional requirements to initiation criteria |
| | For diagnosis of PsA: inadequate response, intolerance or contraindication to a 3-month trial of at least ONE conventional DMARD |
| | For diagnosis of PsO: inadequate response, intolerance, or contraindication to either conventional systemic therapy or phototherapy and if the patient is intolerant or contraindicated to either therapy then the other treatment option needs to be tried |
| | For diagnosis of CD: inadequate response, intolerance or contraindication to at least ONE conventional therapy option and prescriber will initiate dosing of patient with one infusion |
| | Addition of Appendix 1 & 2 |
| September 2018 | Annual editorial review and reference update |
| September 2019 | Annual review |
| November 2019 | Addition of indication: ulcerative colitis. Revised initial dosing requirements for CD |
| December 2019 | Annual review. Addition of requirement to trial preferred product |
| August 2020 | Revised age requirement for plaque psoriasis from 12 and older to 6 and older. Also revised the dosage questions for plaque psoriasis. Clarifying language added to pharmacy benefit |
| September 2020 | Annual review |
| December 2020 | Annual editorial review. Revised requirements to t/f preferred products to apply to Blue Focus patients only. Added PA quantity limits |
| February 2021 | Revised psoriatic arthritis dosing requirement and quantity limits chart |
| March 2021 | Annual editorial review and reference update. Revised background and summary sections. Clarification added to the t/f, intolerance, C/I to preferred products requirement indicating that it only applies to claims adjudicated through the pharmacy benefit. Appendix 1 updated. |
| June 2021 | Annual review |
| March 2022 | Added Conventional Therapy Options for UC chart under Appendix 2 |
| June 2022 | Annual review |
| August 2022 | Per PI update, changed PsA age to 6 and older from 18 and older and updated PsA dosing agreements |
| September 2022 | Annual review |
| December 2022 | Annual review and reference update |
| September 2023 | Annual review |
| March 2024 | Annual editorial review. Revised FDA dosing language |
| May 2024 | Addition of biosimilar Selarsdi |
| June 2024 | Annual review and reference update |
| July 2024 | Addition of biosimilar Pyzchiva |

5.90.004

| | | | |
|--------------------|--------------------|------------------------------|-------------------|
| Section: | Prescription Drugs | Effective Date: | April 1, 2026 |
| Subsection: | Biologicals | Original Policy Date: | November 15, 2013 |
| Subject: | Stelara | Page: | 14 of 17 |

| | |
|----------------|---|
| September 2024 | Annual review |
| January 2025 | Addition of biosimilars Wezlana, Yesintek, and Steqeyma |
| March 2025 | Annual review and reference update |
| April 2025 | Addition of biosimilars Imuldosa and Otulfi |
| June 2025 | Annual review |
| July 2025 | Addition of biosimilar Starjemza |
| September 2025 | Annual review |
| December 2025 | Annual review. Added documentation requirement for non-preferred medications. Removed Blue Focus t/f requirements for preferred medications. Added Appendix 3 |
| March 2026 | Annual review and reference update |

[Keywords](#)

This policy was approved by the FEP® Pharmacy and Medical Policy Committee on March 6, 2026 and is effective on April 1, 2026.

| | | | |
|--------------------|--------------------|------------------------------|-------------------|
| Section: | Prescription Drugs | Effective Date: | April 1, 2026 |
| Subsection: | Biologicals | Original Policy Date: | November 15, 2013 |
| Subject: | Stelara | Page: | 15 of 17 |

Appendix 1 - List of DMARDs

Conventional disease-modifying antirheumatic drugs (DMARDs)

| Generic Name | Brand Name |
|--------------------|-----------------------------|
| azathioprine | Azasan, Imuran |
| cyclophosphamide | Cytoxan |
| cyclosporine | Neoral, Gengraf, Sandimmune |
| hydroxychloroquine | Plaquenil |
| leflunomide | Arava |
| methotrexate | Rheumatrex, Trexall |
| mycophenolate | Cellcept |
| sulfasalazine | Azulfidine, Sulfazine |

Biological disease-modifying antirheumatic drugs (DMARDs)

| Generic Name | Brand Name |
|--------------------|----------------------|
| abatacept | Orencia |
| adalimumab | Humira |
| anakinra | Kineret |
| bimekizumab-bkzx | Bimzelx |
| brodalumab | Siliq |
| certolizumab | Cimzia |
| etanercept | Enbrel |
| golimumab | Simponi/Simponi Aria |
| guselkumab | Tremfya |
| infliximab | Remicade |
| infliximab-dyyb | Zymfentra |
| ixekizumab | Taltz |
| risankizumab-rzaa | Skyrizi |
| rituximab | Rituxan |
| sarilumab | Kevzara |
| secukinumab | Cosentyx |
| spesolimab-sbzo | Spevigo |
| tildrakizumab-asmn | Ilumya |
| tocilizumab | Actemra |
| ustekinumab | Stelara |
| vedolizumab | Entyvio |

Targeted synthetic disease-modifying antirheumatic drugs (DMARDs)

| Generic Name | Brand Name |
|--------------|------------|
|--------------|------------|

Section: Prescription Drugs

Effective Date: April 1, 2026

Subsection: Biologicals

Original Policy Date: November 15, 2013

Subject: Stelara

Page: 16 of 17

| | |
|-----------------|------------|
| apremilast | Otezla |
| baricitinib | Olumiant |
| deucravacitinib | Sotyktu |
| tofacitinib | Xeljanz/XR |
| upadactinib | Rinvoq |

Appendix 2 – List of Conventional Therapies

| Conventional Therapy Options for CD |
|---|
| 1. Mild to moderate disease – induction of remission: a. Oral budesonide, oral mesalamine b. Alternatives: metronidazole, ciprofloxacin |
| 2. Mild to moderate disease – maintenance of remission: a. Azathioprine, mercaptopurine b. Alternatives: oral budesonide, methotrexate intramuscularly (IM) |
| 3. Moderate to severe disease – induction of remission: a. Prednisone, methylprednisolone intravenously (IV) b. Alternatives: methotrexate IM |
| 4. Moderate to severe disease – maintenance of remission: a. Azathioprine, mercaptopurine b. Alternative: methotrexate IM |
| 5. Perianal and fistulizing disease – induction of remission c. Metronidazole ± ciprofloxacin |
| 6. Perianal and fistulizing disease – maintenance of remission d. Azathioprine, mercaptopurine e. Alternative: methotrexate IM |

| Conventional Therapy Options for UC |
|---|
| 1. Mild to moderate disease – induction of remission: a. Oral mesalamine (e.g., Asacol, Lialda, Pentasa), balsalazide, olsalazine b. Rectal mesalamine (e.g., Canasa, Rowasa) c. Rectal hydrocortisone (e.g., Colocort, Cortifoam) d. Alternatives: prednisone, azathioprine, mercaptopurine, sulfasalazine |
| 2. Mild to moderate disease – maintenance of remission: a. Oral mesalamine, balsalazide, olsalazine, rectal mesalamine b. Alternatives: azathioprine, mercaptopurine, sulfasalazine |
| 3. Severe disease – induction of remission: a. Prednisone, hydrocortisone IV, methylprednisolone IV b. Alternatives: cyclosporine IV, tacrolimus, sulfasalazine |
| 4. Severe disease – maintenance of remission: a. Azathioprine, mercaptopurine b. Alternative: sulfasalazine |

| | | | |
|--------------------|--------------------|------------------------------|-------------------|
| Section: | Prescription Drugs | Effective Date: | April 1, 2026 |
| Subsection: | Biologicals | Original Policy Date: | November 15, 2013 |
| Subject: | Stelara | Page: | 17 of 17 |

- | |
|--|
| <p>5. Pouchitis:</p> <ul style="list-style-type: none">a. Metronidazole, ciprofloxacinb. Alternative: rectal mesalamine |
|--|

Appendix 3 - List of Preferred Products

List of preferred products:

https://info.caremark.com/content/dam/enterprise/caremark/microsites/dig/pdfs/pa-fep/fep-misc/FEP_IndicationMedChx.pdf

Refer to formulary documents for confirmation of coverage:

<https://www.fepblue.org/pharmacy/prescriptions#drug-lists>