

Federal Employee Program® 1310 G Street, N.W. Washington, D.C. 20005 202.942.1000 Fax 202.942.1125

5.01.018

Section: Prescription Drugs Effective Date: July 1, 2023

Subsection: Anti-infective Agents Original Policy Date: July 1, 2001

Subject: Itraconazole Page: 1 of 5

Last Review Date: June 15, 2023

Itraconazole

Description

Sporanox (itraconazole); Tolsura* (itraconazole)

*Prior authorization for the brand formulation applies only to formulary exceptions due to being a non-covered medication.

Background

Itraconazole is an oral azole antifungal agent indicated for the treatment of blastomycosis, histoplasmosis, aspergillosis, onychomycosis and oropharyngeal or esophageal candidiasis. FDA-approved indications vary by dosage form, and dosage forms are not interchangeable. Itraconazole works by inhibiting the production of ergosterol (principal sterol in fungal cell membrane) and inhibiting cell membrane formation (1).

Regulatory Status

FDA-approved indications: Sporanox oral solution is indicated for the treatment of oropharyngeal and esophageal candidiasis. Only the oral solution has demonstrated effective for oral and/or esophageal candidiasis (2).

Sporanox capsules and Tolsura capsules are indicated for the treatment of blastomycosis (pulmonary and extrapulmonary), histoplasmosis (including chronic cavitary pulmonary disease and disseminated, non-meningeal histoplasmosis), and aspergillosis (pulmonary and extrapulmonary) in patients who are intolerant of or who are refractory to amphotericin B therapy (2-3).

Sporanox capsules are also indicated for the treatment of onychomycosis in non-immunocompromised patients (3).

5.01.018

Section: Prescription Drugs Effective Date: July 1, 2023
Subsection: Anti-infective Agents Original Policy Date: July 1, 2001

Subject: Itraconazole Page: 2 of 5

Tolsura Limitations of Use: (4)

Tolsura is not indicated for the treatment of onychomycosis and it is not interchangeable or substitutable with other itraconazole products.

Itraconazole has a boxed warning for congestive heart failure, cardiac effects and drug interactions. Do not administer itraconazole in patients with evidence of ventricular dysfunction, such as congestive heart failure (CHF) or a history of CHF. Coadministration of a number of CYP3A4 substrates are contraindicated with itraconazole (2-4).

Itraconazole has warnings regarding hepatic effects, cardiac dysrhythmias, cardiac disease, interaction potential, interchangeability, hydroxypropyl-β-cyclodextrin, and treatment of severely neutropenic patients (2-4)

There are three major forms of aspergillosis: invasive, saprophytic, and allergic. The Infectious Disease Society of America (IDSA) recommends the use of itraconazole and corticosteroids for the treatment of allergic bronchopulmonary aspergillosis (5).

The safety and efficacy of itraconazole in patients less than 18 years of age have not been established (2-4).

Related policies

Cresemba, Ketoconazole, Vfend

Policy

This policy statement applies to clinical review performed for pre-service (Prior Approval, Precertification, Advanced Benefit Determination, etc.) and/or post-service claims.

Itraconazole may be considered **medically necessary** if the conditions indicated below are met.

Itraconazole may be considered **investigational** for all other indications.

Prior-Approval Requirements

Age 18 years of age or older

Diagnoses

Patient must have **ONE** of the following

Section:Prescription DrugsEffective Date:July 1, 2023Subsection:Anti-infective AgentsOriginal Policy Date:July 1, 2001

Subject: Itraconazole Page: 3 of 5

Sporanox oral solution ONLY

- 1. Candidiasis
 - a. Must be unresponsive or refractory to fluconazole
 - b. Includes oropharyngeal and esophageal candidiasis

Sporanox capsules ONLY

- 1. Onychomycosis
 - a. Not immunocompromised
 - b. Not have evidence of ventricular dysfunction, such as congestive heart failure or a history of CHF

Sporanox capsules and Tolsura capsules

- 1. Aspergillosis invasive or saprophytic
 - a. Must be refractory or intolerant to amphotericin B
- 2. Aspergillosis allergic bronchopulmonary
- 3. Blastomycosis
- 4. Coccidioidomycosis
- 5. Histoplasmosis
- 6. Sporotrichosis

AND ALL of the following for ALL diagnoses:

a. Prescriber agrees to monitor for QTc prolongation

Prior - Approval Renewal Requirements

Same as above

Policy Guidelines

Pre - PA Allowance

None

Prior - Approval Limits

Duration 6 months for a diagnosis of Onychomycosis or Candidiasis

1 year for all other diagnoses

5.01.018

Section: Prescription Drugs Effective Date: July 1, 2023
Subsection: Anti-infective Agents Original Policy Date: July 1, 2001

Subject: Itraconazole Page: 4 of 5

Prior - Approval Renewal Limits

Same as above

Rationale

Summary

Itraconazole is an oral azole antifungal agent indicated for the treatment of blastomycosis, histoplasmosis, aspergillosis, onychomycosis and oropharyngeal or esophageal candidiasis. FDA-approved indications vary by dosage form, and dosage forms are not interchangeable. Itraconazole works by inhibiting the production of ergosterol (principal sterol in fungal cell membrane) and inhibiting cell membrane formation. Itraconazole has a boxed warning for congestive heart failure, cardiac effects, and drug interactions. Itraconazole has warnings regarding hepatic effects, interchangeability, hydroxypropyl- β -cyclodextrin, and treatment of severely neutropenic patients. The safety and efficacy of itraconazole in patients less than 18 years of age have not been established (1-4).

Prior approval is required to ensure the safe, clinically appropriate, and cost-effective use of Itraconazole while maintaining optimal therapeutic outcomes.

References

- 1. Itraconazole. Clinical Pharmacology [database online]. Tampa, FL: Elsevier; Revision year 2018. Available from: http://www.clinicalkey.com.
- 2. Sporanox oral solution [package insert]. Titusville, NJ: Janssen Pharmaceuticals; September 2020.
- 3. Sporanox capsules [package insert]. Titusville, NJ: Janssen Pharmaceuticals; December 2019.
- 4. Tolsura [package insert]. Greenville, NC: Mayne Pharma International Pty Ltd; April 2022.
- 5. Patterson TF, Thompson GR, Denning DW, et al. Practice Guidelines for the Diagnosis and Management of Aspergillosis: 2016 Update by the Infectious Diseases Society of America. *Clinical Infectious Diseases*. 2016; 63:112-146.

Policy History	
Date	Action
December 2012	Annual editorial review and reference update.
March 2013	Addition of age and contraindication for CHF
	Addition of Onmel as a line extension
September 2014	Annual editorial review and reference update

5.01.018

Section: Prescription Drugs Effective Date: July 1, 2023
Subsection: Anti-infective Agents Original Policy Date: July 1, 2001

Subject: Itraconazole Page: 5 of 5

September 2015 Annual editorial review and reference update. Addition of identified types of

aspergillosis- allergic bronchopulmonary and aspergillosis - invasive or

saprophytic

March 2016 Annual editorial review

Policy number changed from 5.03.18 to 5.01.18

December 2017 Annual editorial review and reference update

November 2018 Annual editorial review

January 2019 Addition of Tolsura. Policy renamed Itraconazole

March 2019 Annual review

June 2019 Annual review. Added requirement to monitor for QTc prolongation per

SME

September 2019 Removed diagnosis of other fungal infection if patient is diabetic or

immune compromised

December 2019 Annual review. Moved Tolsura to MFE with PA only.

December 2020 Annual review and reference update

June 2021 Annual editorial review and reference update. Removed Onmel from policy

due to being discontinued. Criteria revised so that Sporanox Oral Solution

is the only formulation approved for candidiasis to align with PI

December 2021 Annual review

June 2022 Annual review and reference update

December 2022 Annual review and reference update. Changed policy number to 5.01.018

June 2023 Annual review

Keywords

This policy was approved by the FEP® Pharmacy and Medical Policy Committee on June 15, 2023 and is effective on July 1, 2023.